

CHOICE WITH ENDOGENOUS CATEGORIZATION*

ANDREW ELLIS[†] AND YUSUFCAN MASATLIOGLU[§]

ABSTRACT. We propose a novel categorical thinking model (CTM) where the framing of the decision problem affects how the agent categorizes each product, and the product's category affects her evaluation of the product. We show that a number of prominent models of salience, status quo bias, loss-aversion, inequality aversion, and present bias all fit under the umbrella of CTM. This suggests categorization as an underlying mechanism for key departures from the neoclassical model of choice and an account for diverse sets of evidence that are anomalous from its perspective. We specialize CTM to provide a behavioral foundation for the salient thinking model of Bordalo et al. [2013], highlighting its strong predictions and distinctions from other existing models.

Date: March, 2020.

*We thank David Dillenberger, Erik Eyster, Nicola Gennaioli, Matt Levy, Collin Raymond, the anonymous referees, Andrei Shleifer, Rani Spiegler, Tomasz Strzalecki, and conference/seminar participants at BRIC 2017, CETC 2017, SAET 2017, Lisbon Meetings 2017, ESSET 2019, Brown, UPenn, Pompeu Fabra, and Harvard for helpful comments and discussions. This project began at ESSET Gerzensee, whose hospitality is gratefully acknowledged.

[†] Department of Economics, London School of Economics, Houghton Street, London, WC2A 2AE. Email: a.ellis@lse.ac.uk.

[§] University of Maryland, 3147E Tydings Hall, 7343 Preinkert Dr., College Park, MD 20742. E-mail: yusufcan@umd.edu.

1. INTRODUCTION

Psychologists have long held that knowledge about our environment is organized into categories, and that this categorization plays a key role in decision making. Categorization has been used by both humans and animals for thousands of years. As Ashby & Maddox [2005] write, “*All organisms assign objects and events in the environment to separate classes or categories... Any species lacking this ability would quickly become extinct.*”

Categorization plays a key role in a number of important anomalies for the neo-classical model of choice. Attributes categorized as losses get higher weight relative to those categorized as gains [Tversky & Kahneman, 1991]. An object’s most salient attribute plays a disproportionate role in the agent’s subsequent evaluation [Bordalo et al., 2013]. Subjects avoid objects they categorize as not-obviously-better-than the status-quo [Masatlioglu & Ok, 2005]. Agents are less patient when deciding between dated rewards in the short-term than in the long-term [Strotz, 1955]. Allocations among members of society are evaluated according to whether inequities are advantageous or disadvantageous [Fehr & Schmidt, 1999].

This paper proposes and axiomatizes a simple model of the role that categorization plays in economic decisions. In the Categorical Thinking Model (CTM), a decision maker (DM) first groups objects together into categories, consciously or unconsciously, then evaluates each object through the lens of the category to which it belongs. The model has two key features motivated by psychological evidence. First, categorization is context-dependent, as summarized by a reference point that may depend on the choice set. Second, how an object is categorized affects its valuation. Prominent models of loss-aversion, salience, status quo bias, present bias, and inequality aversion all fit under the umbrella of CTM. Hence, CTM suggests categorization as an underlying explanation for many key departures from the neoclassical model in many different decision-making environments.

To make our results comparable with previous work, we begin by assuming that a family of reference-dependent preference relations describe the DM’s choices for each reference point. Each alternative has a pair of observed attributes, such as price and quality, height and weight, or size and timing of a reward. In CTM, the context in which the decision takes place determines a reference point, which in turn divides the alternatives into categories. Each category has its own utility function, and within a given category, the DM evaluates the options according to it. Hence, the DM makes different trade-offs between the attributes when they are differentially categorized. We show that the DM conforms to CTM if and only if she behaves as a standard DM when comparing objects categorized the same way. That is, her choices satisfy some standard axioms, such as acyclicity, and do not depend on the reference point when restricted to alternatives that belong to the same category.

CTM is a parsimonious approach to incorporating psychological evidence into economics. Psychological factors determine how each alternative is perceived, which CTM captures through different categories. Moreover, they predict how being categorized in a particular way affects the DM’s choice, which CTM captures through the category’s utility function. For instance, salience and loss-aversion make distinct predictions about when a DM puts higher weight on a dimension. The most salient attribute gets more weight, as does an attribute classified as a loss. Our result shows that CTM closes the model by requiring that the DM acts consistently within the alternatives categorized the same way.

Despite its generality, CTM makes testable predictions and excludes certain types of modeling choices. For instance, a number of models capture salience effects, including the salient thinking model [Bordalo et al., 2013] (BGS), Kőszegi & Szeidl [2013], Bhatia & Golman [2013], Gabaix [2014], and Bushong et al. [2015]. Of these models, only BGS is a CTM. In other words, even the most general version of BGS excludes these models, so BGS offers a different method of modeling salience. Our results highlight trade-offs between the different modeling approaches. For instance, BGS maintains a stronger consistency condition across reference points than does the

constant loss aversion of [Tversky & Kahneman, 1991], but the latter, unlike BGS, satisfies Monotonicity across regions.

We then provide the first complete characterization of the observable choice behavior equivalent to the BGS model, clarifying and identifying the nature of the assumptions used in the model. The first crucial step towards understanding the model is getting a handle on its novel *salience function* that determines which attribute stands out for a given reference point. We study the salience function based on a simple observation: while it influences which attribute is salient, the weight given to each attribute is independent of its magnitude. This makes BGS a special case of CTM, so our earlier results allow a characterization.

One key feature of BGS is that the reference point is endogenously determined by the set of available options. Since the salience of each alternative depends on the reference point, varying the budget set affects the salience of, and so the DM’s evaluation of, a given alternative. Our final contribution addresses this challenge by extending our characterization of CTM to the setting where the reference point is endogenous. Our primitive is a choice correspondence describing the DM’s choices. The menu maps to a reference point, such as the average level of each attribute over alternatives in the set. As long as the reference point varies systematically with the choice problem, we characterize the properties of the choice correspondence equivalent to CTM. Specifically, we show that if the DM’s choices obey the natural analogs of our earlier axioms, then CTM rationalizes her behavior. We apply it to provide a completely endogenous characterization of the BGS function.

The paper proceeds as follows. The next subsection provides a brief overview of the relevant psychology literature on categorization. Section 2 introduces CTM and discusses the models covered under its umbrella. Section 3 axiomatizes CTM and compares and contrasts the models of riskless choice discussed in Section 2. Section 4

contains our analysis of the salient thinking model. Section 5 introduces the endogenous reference point setting, and applies our axiomatizations of CTM to it. Section 6 concludes with a discussion of related literature.

1.1. Psychology of Categorization. There is a long literature in psychology and marketing discussing categorization. Recent review articles include Ashby & Maddox [2005], Loken [2006], Loken et al. [2008] and Cosmides & Tooby [2013]. Much of the literature focuses on how categories are formed, and when new alternatives are added into existing categories. CTM relies on several properties documented by this literature.

First, categories are context dependent. Tversky [1977], Tversky & Gati [1978] present evidence that replacing one item in a set of objects can drastically alter how people categorize the remaining objects. Tversky & Gati [1978] argue that categorization “*is generally not invariant with respect to changes in context or frame of reference.*” For example, they show that subjects put East Germany and West Germany into the same category when the salient feature is geography or cultural background, but categorize the two differently if political system is salient. Similarly, Choi & Kim [2016] posit that depending on the context an Apple Watch can be categorized as a tech product, a fashion product, a fitness product, or a simple watch. Ratneshwar & Shocker [1991] show that subjects categorize ice cream and cookies together in terms of similarity (e.g. they are both desserts), but categorize ice cream and hot dogs together in terms of usage benefit (e.g. both are good snacks to have at the pool). Stewart et al. [2002] present evidence that relative magnitude information, derived from a comparison of the reference point, is used in categorization of sounds.

Second, how an object is categorized affects its final valuation. In a classic series of experiments, Rosch [1975] shows that differently categorized but physically identical stimuli are perceptually encoded as distinct objects. Wanke et al. [1999] demonstrate that “wine” is evaluated more positively when categorized with “lobster” than with “cigarettes.” Mogilner et al. [2008] show that categorizing goods differently resulted

in different reported satisfaction. Chernev [2011] shows that bundling a healthy food item with a junk food item reduced the reported caloric content beyond that of the junk food alone.

Finally, categories take the form of regions in the alternative space. This tracks very closely with the decision bound theory in the psychology. As Ashby & Maddox [2005, p. 152] describe, it posits that the subject “*partition[s] the stimulus space into response regions... determines which region the percept is in, and then emits the associated response.*” Ashby & Gott [1988] show it can accommodate examples incompatible with other theories of category formation, such as prototype theory. Moreover, there is substantial experimental support for it, including Ashby & Waldron [1999], Anderson [1991], Love et al. [2004].

2. MODEL

To aid in comparison with the existing literature and to separate the effects of reference point formation, we follow Tversky & Kahneman [1991] by taking as given a family of reference-dependent preference relations. We assume that the space of alternatives is $X = \mathbb{R}_{++}^n$, focusing on $n = 2$ when not otherwise noted.¹ We often use the convention of writing x as (x_i, x_{-i}) with x_{-i} denoting the components of x different for i . The next subsections explore three different interpretations of X in different contexts: as a riskless object with different attributes, as a dated reward or consumption stream, and as an allocation of consumption across individuals. For each reference point $r \in X$, the DM maximizes a complete and transitive preference relation, denoted by \succsim_r , over X . As usual, \succ_r denotes strict preference and \sim_r indifference. The primitive of the model is a family of such preferences indexed by the set of reference points, $\{\succsim_r\}_{r \in X}$. In this section, we assume that the reference point is exogenously

¹ We note when there is a distinction between general n and $n = 2$. Theorem 5 and the results that rely on it use the full structure of \mathbb{R}_{++}^n . The remaining results all generalize to any X that is a finite Cartesian product of open, linearly ordered, separable, connected sets endowed with the order topology, where X itself has the product topology.

given. We relax this assumption in Section 5 to allow endogenous reference point formation.

2.1. Categorical Thinking Model. The first ingredient of the model is a mapping from the reference r to categories. Each category corresponds to a different psychological treatment and changes as the reference changes. We allow the categories to have a very general structure.

Definition 1. A vector-valued function $\mathcal{K} = (K^1, K^2, \dots, K^m)$ is a *category function* if each $K^k : X \rightarrow 2^X$ satisfies the following properties:

- (1) $K^k(r)$ is a non-empty, regular open set, and $cl(K^k(r))$ is connected,²
- (2) $\bigcup_{k=1}^m K^k(r)$ is dense,
- (3) $K^k(r) \cap K^l(r) = \emptyset$ for all $k \neq l$, and
- (4) $K^k(\cdot)$ is continuous.³

Categories arise from the psychology of the phenomenon to be modeled. For CTM to be applicable, the psychology must make an unambiguous prediction about which alternatives are affected. For instance, with gain-loss utility, alternatives that dominate the reference point are treated differently than those better in only one dimension. Similarly, with present-bias, alternatives that pay-off sooner than the reference are categorized together. While we take the categories as given, if the psychology only makes partial predictions, then the categorization of other alternatives can often be inferred from choice. Proposition 1 does so for the salient thinking model.

We interpret the properties as follows. Every category contains some alternative for every reference point. If a particular product, say x , belongs to the category k , then so do all products that are close enough to x . There is a path that stays within the category between any two points, so categories cannot be the union of “islands.” Almost every alternative is in at least one category, and none are in two categories. Finally, if the reference point does not change too much, then neither do the categories.

²Recall that a set A is regular open if $A = int(cl(A))$.

³That is, each K^k is both upper and lower hemicontinuous when viewed as a correspondence.

The consumer values each good in a way that depends not only on alternative of a product, as in the standard neoclassical model, but also on the category to which the product belongs. When alternatives x and y are both categorized in category k , the *category utility function* $U^k : X \rightarrow \mathbb{R}$ represents the DM's choices. That is, she prefers x to y if and only if $U^k(x) \geq U^k(y)$. We focus on the effect of categorization on distorting trade-offs, so we require that a category utility function is *additively separable* and *monotonic*: $U^k(x) = \sum_{i=1}^n U_i^k(x_i)$ where each $U_i^k(\cdot)$ is strictly monotone and continuous.⁴ The utility index U_i^k represents the DM's preferences over dimension i when an alternative belongs to the category k .

When alternatives belong to different categories, the reference point may affect the DM's choice. If the alternative x lies in the category k when the reference is r , that is, $x \in K^k(r)$, then the value of consumption x is represented by $U^k(x|r)$. However, the reference does not affect the utility trade-off within a category. To capture this, we require that $U^k(\cdot|r)$ agrees with U^k , in the sense that it is an increasing transformation thereof. Then, $U^k(x|r) \geq U^k(y|r)$ if and only if $U^k(x|r') \geq U^k(y|r')$ for any references $r, r' \in X$. We can now formally define the model as follows.

Definition 2. The family $\{\succsim_r\}_{r \in X}$ conforms to the *Categorical Thinking Model (CTM)* under category function $\mathcal{K} = (K^1, K^2, \dots, K^m)$ if for each category k there is a *category utility function* U^k so that when $x \in K^k(r)$ and $y \in K^l(r)$ for some r

$$x \succsim_r y \iff U^k(x|r) \geq U^l(y|r)$$

and $U^k(\cdot|r)$ is an increasing transformation of $U^k(\cdot)$ for each $r \in X$ and category k .

A CTM is *increasing* if U_i^k is increasing in x_i for every category k and dimension i . We also consider two sub-classes: A CTM is *affine* if $U^k(\cdot|r)$ an affine transformation of U^k for each r . A CTM is *strong* if $U^k(\cdot|r) = U^k(\cdot)$ for each r . Most of the models we discuss below are affine CTM, and those of riskless consumer choice are all increasing.

⁴That is, U_i^k is either strictly increasing on \mathbb{R}_+ or strictly decreasing on \mathbb{R}_+ .

2.2. Riskless Consumer Choice. In this subsection, we consider our primary application: riskless consumer choice. The four models are introduced formally, and each is shown to be CTM. Figure 1 plots their indifference curves and categories, with darker lines indicating higher utility.




FIGURE 1. CTM for Riskless Choice

SALIENT THINKING MODEL (BGS): Bordalo et al. [2013] propose an intuitive and descriptive behavioral model based on salience. In the model, an attribute receives more weight when it is salient than when it is not. The magnitude of salience is determined by a *salience function*, $\sigma := \mathbb{R}_{++} \times \mathbb{R}_{++} \rightarrow \mathbb{R}_+$. Given a reference (r_1, r_2) , attribute 1 is salient for good x if $\sigma(x_1, r_1) > \sigma(x_2, r_2)$, and attribute 2 is salient for good x if $\sigma(x_1, r_1) < \sigma(x_2, r_2)$.⁵ That is, the salient attribute is the one that differs the most from the reference according to the salience function.

⁵We describe the properties of σ more fully in Section 4.

Definition 3. The family $\{\succsim_r\}_{r \in X}$ has a BGS $(\sigma; w_1, w_2, u_1, u_2)$ representation if each \succsim_r is represented by

$$(1) \quad V_{BGS}(x|r) = \begin{cases} w_1^1 u_1(x_1) + w_2^1 u_2(x_2) & \text{if } \sigma(x_1, r_1) > \sigma(x_2, r_2) \\ w_1^2 u_1(x_1) + w_2^2 u_2(x_2) & \text{if } \sigma(x_2, r_2) > \sigma(x_1, r_1) \end{cases}$$

for a salience function σ , strictly positive weights with $\frac{w_1^1}{w_2^1} > \frac{w_1^2}{w_2^2}$, and each u_i strictly increasing.

To illustrate this model, consider the salience function proposed by BGS:

$$\sigma(x_k, r_k) = \frac{|x_k - r_k|}{x_k + r_k}.$$

Based on it, the left-upper panel in Figure 1 illustrates BGS. There are two categories: those that are 1-salient, i.e. $\sigma(x_1, r_1) > \sigma(x_2, r_2)$, and those that are 2-salient, i.e. $\sigma(x_2, r_2) > \sigma(x_1, r_1)$. To visualize them, note that the entire product space is divided into four distinct areas by the two dashed curves that intersect at the reference point. The areas lying the north and south of the reference point are categorized as the 2-salient products. Similarly, 1-salient products lie east and west of the reference point. The figure incorporates indifference curves as well, holding fixed the reference point. There are two potential sets of indifference curves, illustrated by dotted lines. Depending on the category, one of the two is utilized to determine the DM's choice. When attribute 1 is salient, the steeper one becomes the indifference curve since it puts higher weight on the first attribute. Conversely, the flatter one is the indifference curves when attribute 2 is salient. We draw two different indifference curves, where the darker color corresponds to higher utility.

CONSTANT LOSS AVERSION MODEL (TK): Tversky & Kahneman [1991] provides foundations for a reference-dependent model that extends Prospect Theory to riskless consumption bundles. Each is evaluated relative to reference point r , and losses loom larger than gains. In the absence of losses, the DM values each alternative with an additive utility function, $u(x_1) - u(r_1) + v(x_2) - v(r_2)$, which attaches equal weight

to each attribute. If she experiences a loss in attribute i , then she inflates the weight attached to that attribute by $\lambda_i > 1$. There are four different categories in the TK formulation: (i) gain in both dimensions, (ii) gain in the first dimension and loss in the second dimension, (iii) loss in the first dimension and gain in the second dimension, and (iv) loss in both dimensions (see the right-upper panel in Figure 1). We model this as $\mathcal{K}^{GL} = (K^1, K^2, K^3, K^4)$ where $K^1(r) = \{x : x \gg r\}$, $K^2(r) = \{x : x_1 < r_1 \text{ and } x_2 > r_2\}$, $K^3(r) = \{x : x_1 > r_1 \text{ and } x_2 < r_2\}$, and $K^4(r) = \{x : x \ll r\}$; call this the *gain-loss category function*. Then, the utility function is

$$V_{TK}(x|r) = \begin{cases} u_1(x_1) - u_1(r_1) + u_2(x_2) - u_2(r_2) & \text{if } x \in K^1(r) \\ \lambda_1(u_1(x_1) - u_1(r_1)) + u_2(x_2) - u_2(r_2) & \text{if } x \in K^2(r) \\ u_1(x_1) - u_1(r_1) + \lambda_2(u_2(x_2) - u_2(r_2)) & \text{if } x \in K^3(r) \\ \lambda_1(u_1(x_1) - u_1(r_1)) + \lambda_2(u_2(x_2) - u_2(r_2)) & \text{if } x \in K^4(r) \end{cases}$$

where $\lambda_1, \lambda_2 > 0$ (> 1 if loss averse) and each u_i strictly increasing. TK is a special case of affine CTM with four categories defined by a gain-loss category function.

STATUS QUO BIAS MODEL (MO): Masatlioglu & Ok [2005] model individuals who experience some form of psychological discomfort when they have to abandon their status quo option. This discomfort imposes an additional utility cost. Of course, if an alternative is unambiguously superior to the status quo, the DM does not feel any psychological discomfort to forgo the status quo; in such cases there will be no cost. Formally, $Q(r)$ is a closed set denoting the alternatives that are unambiguously superior to the default option r (see the left-bottom panel of Figure 1). If an alternative does not belong to this set, then the DM pays a cost $c(r) > 0$, which may depend on the reference point, to move away from the status quo. In this model, there are two categories $\mathcal{K}^{MO} = (K^1, K^2)$ where $K^1(r) = \{x | x \in \text{int}(Q(r))\}$ and $K^2(r) = \{x | x \notin Q(r)\}$. For any $x \neq r$, we have

$$V_{MO}(x|r) = \begin{cases} u_1(x_1) + u_2(x_2) & \text{if } x \in K^1(r) \\ u_1(x_1) + u_2(x_2) - c(r) & \text{if } x \in K^2(r) \end{cases}.$$

This is an example of an affine CTM for general c , and a strong CTM when $c(r)$ is constant.

PROTOTYPE THEORY (PT): Prototype theory was first proposed by Posner & Keele [1970]. According to it, each category is associated with a prototype, its “most typical” member. Initial categorization is determined by comparing each product to each prototype. We now formalize this idea and show that this is CTM.

There are m prototypes, p^1, \dots, p^m . The DM categorizes alternatives according to how similar they are to a given prototype. Then, category $K^i(r)$ is the set of alternatives categorized as most similar to exemplar p^i . Similarity may depend on the reference. There is a family of metrics indexed by r so that $d_r(x, y)$ indicates how far away the DM perceives x to be from y given reference r . Formally, $\mathcal{K}^P = (K^1, \dots, K^m)$ where

$$K^i(r) = \{x : i = \arg \min_j d_r(p^j, x)\}.$$

and the DM evaluates alternatives in category i according to

$$V_{PT}^i(x|r) = U(p^i) + \lambda_1^i(x_1 - p_1^i) + \lambda_2^i(x_2 - p_2^i) \text{ if } x \in K^i(r)$$

where $U(\cdot)$ is a hedonic utility function and $\lambda_j^i > 0$. A particularly interesting specification is where $\lambda_j^i = \frac{\partial}{\partial p_j^i} U(p^i)$. Then, the DM approximates the utility of x according to a first-order Taylor expansion around the prototype most similar to it (see the right-bottom panel of Figure 1).⁶ This is an example of a Strong CTM.

2.3. Time Preferences. We apply our model to choices of dated rewards. The pair (x, t) represents a payment of x at time t . Motivated by present bias, we propose a model where the DM divides time periods according to short term and long term. Given a reference $r = (r_x, r_t)$, rewards arriving before r_t are perceived as a short-term and after r_t as long-term. Hence

$$K^{short}(r) = \{(x, t) | t < r_t\}$$

⁶In the figure, we use $d_r(p^j, x) = \frac{d(p^j, x)}{d(p^j, r)}$ where d is the Euclidean metric.

and

$$K^{long}(r) = \{(x, t) | t > r_t\}.$$

The utility function is

$$V_{QH}((x, t) | (r_x, r_t)) = \begin{cases} (\beta\delta)^t u(x) & \text{if } (x, t) \in K^{short}(r) \\ \beta^{r_t} \delta^t u(x) & \text{if } (x, t) \in K^{long}(r) \end{cases}$$

where $0 < \delta < 1$ and $0 < \beta \leq 1$. The model is additively separable after taking logs, so it is a special case of CTM. It exhibits present bias when $\beta < 1$: there exist values $y > x > 0$ so that the DM prefers $(x, \tau) \succsim_r (y, \tau + 1)$ if and only if $\tau < r_t - 1$.⁷ Figure 2 plots its indifference curves.




FIGURE 2. CTM for Dated Rewards

2.4. Social preferences. Our final application is to consumption allocations. Alternatives assign consumption to each of n agents, labeled $1, \dots, n$. Dimension 1 corresponds to the DM's own consumption. We consider inequality-averse and social-welfare concerned DMs.

INEQUALITY AVERSION: Fehr & Schmidt [1999] introduce a model of inequality aversion. The DM experiences envy, i.e. she dislikes having a lower allocation than another,

⁷For instance $u(x) = 1$ and $u(y) = (\beta\delta)^{-1}$.

and guilt, i.e. she prefers others not to have less consumption than her. We present a generalization of their model where a reference point affects how much envy or guilt the DM feels. Envy and guilt are generated by the difference between how much better agent i is relative to i 's reference consumption and how much better the DM is relative to her own reference. Some reference-dependence makes sense when considering decisions that impact the social alternative: the DM may not experience much guilt if agent 2's consumption is low in every feasible allocation.

In the *Relative Inequality Aversion (RIA)* model, the DM i feels guilty if her own relative gain $(x_i - r_i)$ is higher than the relative gain of individual j $(x_j - r_j)$. Otherwise, the individual i is jealous of individual j since $x_j - r_j > x_i - r_i$. Given the reference point is r , the value function of the DM equals

$$V_i^{RIA}(x|r) = x_i - \frac{\alpha}{(n-1)} \sum_{j \neq i} \max\{(x_j - r_j) - (x_i - r_i), 0\} \\ - \frac{\beta}{(n-1)} \sum_{j \neq i} \max\{(x_i - r_i) - (x_j - r_j), 0\}$$

Observe that when $r_i = r_j$ for all i and j , the utility function reduces to that of Fehr & Schmidt [1999]. Throughout, we follow them in assuming that $\alpha \geq \beta \geq 0$ and $\beta < 1$

We illustrate that this model is CTM by using the two-person version of it. With two individuals, the category function can be written as $\mathcal{K}^{RIA} = (K_R^1, K_R^2)$ where

$$K_R^1(r) = \{x \in X : x_1 - r_1 > x_2 - r_2\} \text{ and } K_R^2(r) = \{x \in X : x_1 - r_1 < x_2 - r_2\}$$

The set $K_R^j(r)$ contains allocations where individual j gets a relatively better deal than the other. The relative inequality aversion model can be written as

$$V_1^{RIA}(x|r) = \begin{cases} x_1 - \alpha[(x_1 - r_1) - (x_2 - r_2)] & \text{if } x \in K^1(r) \\ x_1 - \beta[(x_2 - r_2) - (x_1 - r_1)] & \text{if } x \in K^2(r) \end{cases}$$

which is an affine CTM.

DISTRIBUTIONAL PREFERENCES: Charness & Rabin [2002] propose a model of social preferences where utility is increasing with the minimum of all individuals' payoffs and




FIGURE 3. Left: Relative Inequality Aversion and Right: Reference-Dependent Distributional Preferences

the total of all individuals' payoffs. The DM i maximizes

$$V_i(x) = (1 - \lambda)x_i + \lambda[\delta \min\{x_1, x_2, \dots, x_n\} + (1 - \delta) \sum_k x_k].$$

The parameter $\delta \in (0, 1)$ measures the degree of concern for helping the worst-off individual (Rawlsian) versus maximizing the total social payoffs (Utilitarian). The parameter $\lambda \in (0, 1)$ measures how the DM balances social welfare with her own utility, where $\lambda = 0$ captures pure self-interest.

We propose a natural extension of their model with an exogenously given reference point. We call this model *Reference-Dependent Distributional Preferences*. That is,

$$V_i^{CR}(x|r) = (1 - \lambda)x_i + \lambda[\delta \min\{x_1 - r_1, x_2 - r_2, \dots, x_n - r_n\} + (1 - \delta) \sum_k x_k]$$

According to this model, each individual cares to maximize the minimum possible relative payoff $x_j - r_j$. Note that if $r_i = r_j$ for all i and j , this model encompasses the model of Charness & Rabin [2002] as a special case.

We show that this model is CTM. To do that, we first define categories for this model. Each category corresponds to the individual who has the worst relative payoff. In this case, $\mathcal{K}^{CR} = (K^1, \dots, K^n)$ where

$$K^j(r) = \{x \in X : (x_j - r_j) \text{ is the minimum of } \{x_1 - r_1, x_2 - r_2, \dots, x_n - r_n\}\},$$

and

$$V_i^{CR}(x|r) = (1 - \lambda)x_i + \lambda[\delta(x_j - r_j) + (1 - \delta) \sum_k x_k] \text{ if } x \in K^j(r),$$

showing that the model is an affine CTM.

3. BEHAVIORAL FOUNDATION FOR CTM

In this section, we provide a set of behavioral postulates characterizing increasing CTM. These postulates represents the key features of the model. We show that they hold if and only if the data is representable by increasing CTM, rendering the model behaviorally testable. In subsequent subsections, we explore the various strengthenings of the model and provide axiomatizations of these as well.

For each category k , define the revealed ranking within that category \succsim^k so that $x \succsim^k y$ if and only if there exists r such that $x, y \in K^k(r)$ and $x \succsim_r y$. The sub-relations \succ^k and \sim^k are defined in the usual way. The ranking \succsim^k captures preference within category k . The following axiom states that the within-category revealed preference has no cycles.

Axiom 1 (Weak Reference Irrelevance). The relation \succsim^k is acyclic. That is, if $x^1 \succsim^k x^2 \succsim^k \dots \succsim^k x^m$, then $x^m \not\succsim^k x^1$.

Weak Reference Irrelevance ensures that the DM reacts consistently to alternatives when they are categorized the same way. That is, the categories reflect the DM's psychological treatment of the alternative. Although she may have choice cycles, these cycles occur only when the context changes how the DM categorizes alternatives. Since \succsim^k is acyclic, we can take its transitive closure to derive full comparisons. Let \succsim^{k*} be its transitive closure, with \succ^{k*} and \sim^{k*} the asymmetric and symmetric parts.

Within a category, preference has an additive structure. The next axiom implies that each \succsim_r satisfies Cancellation when restricted to a given category.

Axiom 2 (Category Cancellation). For all $x_1, y_1, z_1, x_2, y_2, z_2 \in \mathbb{R}_+$, $r \in X$, and category j so that $(x_1, z_2), (z_1, y_2), (z_1, x_2), (y_1, z_2), (x_1, x_2), (y_1, y_2) \in K^j(r)$:

If $(x_1, z_2) \succsim_r (z_1, y_2)$ and $(z_1, x_2) \succsim_r (y_1, z_2)$, then $(x_1, x_2) \succsim_r (y_1, y_2)$.

Category Cancellation adapts the well-known Cancellation axiom to our setting, differing in its requirement that the alternatives belong to the same category. Without the qualifiers on how alternatives are categorized, the axiom is a well-known necessary condition for an additive representation that appears in Krantz et al. [1971] and Tversky & Kahneman [1991], among others. If X has strictly more than two dimensions, then we can replace it with the analog of P2 [Savage, 1954]; see Debreu [1959].⁸

The next axiom requires that Monotonicity holds between objects categorized the same way.

Axiom 3 (Category Monotonicity (CM)). For any $x, y, r \in X$: if $x \geq y$ and $x \neq y$, then $y \not\prec^{k*} x$ for any category k ; in particular, if $x, y \in K^k(r)$, then $x \succ_r y$.

Since both attributes are “goods” as opposed to “bads,” Monotonicity means that if a product x contains more of some or all attributes, but no less of any, than another product y , then x is preferred to y . The postulate requires that choice respects Monotonicity for alternatives within the same category. However, it does not require that this comparison holds when the goods belong to different categories, and we shall see later that salience can distort comparisons enough to cause Monotonicity violations.

Finally, the family of preference relations is suitably continuous.

Axiom 4 (Category Continuity). For any $r \in X$ and any $x \in \bigcup_i K^i(r)$, the sets $UC_j(x) = \{y \in K^j(r) : y \succ_r x\}$ and $LC_j(x) = \{y \in K^j(r) : x \succ_r y\}$ are open.

⁸Formally, for any $x, y, x', y' \in K^k(r)$ and subset of indexes E , if $x_i = x'_i$ and $y_i = y'_i$ for $i \in E$, $x_i = y_i$ and $x'_i = y'_i$ for all $i \notin E$, and $x \succsim_r y$, then $x' \succsim_r y'$. This is implied by Category Monotonicity when $n = 2$, so a stronger condition is necessary.

Moreover, the set

$$\left\{ x \in \bigcup_i K^i(r) : UC_j(x) \cup LC_j(x) = K^j(r) \text{ and } UC_j(x) \neq K^j(r) \text{ and } LC_j(x) \neq K^j(r) \right\}$$

has an empty interior.

Category continuity adapts the usual continuity condition to apply only within a category. It says that when y is preferred to x in a given context and y' is close enough to y , then y' is also preferred to x , provided that y' belongs to the same category as y . The final condition requires that if an alternative x is neither better than everything within category j nor worse than everything within category j , then there exists something in category j that is as good as x , or as good as something arbitrarily close to x . For such an x , the category must intersect almost all indifference curves close to x 's since each category is almost connected.

Finally, we make a structural assumption.

Assumption (Structure). The category function \mathcal{K} is such that for any category k , the following sets are connected: $E^k = \bigcup_{r \in X} K^k(r)$, $\{x \in E^k : x_i = s\}$ for all dimensions i and scalars s , and $\{y \in E^k : x \sim^{k*} y\}$ for all $x \in E^k$.

The Structure Assumption is satisfied all the models we discussed in the previous section. Indeed, $E^k = \mathbb{R}_{++}^n$ for every category in these models. These conditions establish that the objects categorized in the same way have enough topological structure so that “local” properties can be extended to global ones. Chateauneuf & Wakker [1993] show that the structure assumption, applied to a single preference relation and domain, is needed to guarantee that a local additive representation implies a global one.

Theorem 1. *Assume the Structure Assumption holds. The family $\{\succsim_r\}_{r \in X}$ satisfies Weak Reference Irrelevance, Category Cancellation, Category Monotonicity, and Category Continuity for \mathcal{K} if and only if it conforms to increasing CTM under \mathcal{K} .*

Increasing CTM captures the behavior implied by the axioms, so we call Axioms 1-4 the CTM axioms. Taken together, they establish that the DM acts rationally when restricting attention to alternatives categorized in the same way for a given reference point. That is, CTM captures a DM who differs from the neoclassical model only when alternatives are categorized differently. The theorem reveals that a number of other reference dependent models have been studied by the literature fall outside the scope of our analysis. For instance, Bhatia & Golman [2013], Munro & Sugden [2003], the non-constant loss averse version of Tversky & Kahneman [1991], and the continuous version of the salient thinking model (see online appendix of Bordalo et al. [2013]) all violate weak reference irrelevance for any specification of the category function. We provide the details in Appendix A.6.

We provide a brief outline of how the proof works, and all omitted proofs can be found in the appendix. The axioms are sufficient for a “local” additive representation of \succsim_r (and thus \succsim^k) on an open ball around each alternative within category k . The Structure Assumption allows us to apply Theorem 2.2 of Chateauneuf & Wakker [1993] to aggregate the local additive representation of \succsim^k into a global one. To do so, we must establish that the global preference is complete, transitive, monotone, and continuous. We establish these properties for preference within each category by showing that the transitive closure of each \succsim^k is complete and suitably continuous. The remainder of the proof shows that Categorical Continuity allows us to stitch the different within-category representations together into an overall utility function.

3.1. Reweighting. In all of the models discussed in Section 2.2, the DM evaluates the difference between alternatives categorized in the same way similarly. That is, regardless of the category, the DM agrees on how much better a value of x versus y is in dimension i . Categorization affects only how much weight she puts on each dimension. This is captured by the following axiom.

Axiom 5 (Reference Interlocking). For any $a, b, a', b', x', y', x, y \in X$ and categories k, j with $x_{-i} = a_{-i}$, $y_{-i} = b_{-i}$, $x'_{-i} = a'_{-i}$, $y'_{-i} = b'_{-i}$, $x_i = x'_i$, $y_i = y'_i$, $a_i = a'_i$, $b_i = b'_i$: if $x \sim^k y$, $a \succsim^k b$, and $x' \sim^j y'$, then it does not hold that $b' \succ^j a'$.

The term “Reference Interlocking” comes from Tversky & Kahneman [1991]. If each \succsim^k is complete, then their statement of it is equivalent given the other axioms. Roughly, the DM agrees on the difference in utilities along a given dimension regardless of how an alternative is categorized. To interpret, observe that the first pair of comparisons reveals that the difference between a_i and b_i exceeds that between x_i and y_i when the alternatives belong to category k . For alternatives categorized in j , the DM should not reveal the opposite ranking. We defer to the above paper for a detailed discussion.

Theorem 2. *Suppose that $\{\succsim_r\}_{r \in X}$ conforms to increasing CTM under \mathcal{K} and each E^k is connected. For each dimension i , there exist a utility index u_i and a weight $w_i^k > 0$ for each category k so that each category utility U^k is cardinally equivalent to one that maps each $x \in E^k$ to $\sum_i w_i^k u_i(x_i)$ if and only if Reference Interlocking holds.*

All of the models in Section 2.2 satisfy the axiom, and are thus special cases of increasing CTM satisfying Reference Interlocking. For instance, differences in the salient dimension of BGS receive higher weight, but the relative size of two given differences in the same dimension is the same regardless of whether both are salient or both are not. The axiom implies that the utility index within each category must be the same, up to an increasing, affine transformation.

3.2. Behavioral Foundation for Affine CTM. In this section, we explore when an affine CTM exists. That is, when is $U^k(\cdot|r)$ a positive affine transformation of $U^k(\cdot|r')$ for any r, r' ? All of the models from Section 2.2 fall into this class.⁹

⁹For MO, this is true only when $c(r) < \infty$.

Unsurprisingly, the key restriction relative to CTM is that tradeoffs across categories are affine. As is usual, this is captured by a form of linearity, or the “Independence Axiom.” We require it to hold only when alternatives combined belong to the same category, and adjust for the curvature of the utility index.

To state the key axiom, we define an operation \oplus^k along similar lines as Ghirardato et al. [2003]. For $x, y \in \mathbb{R}$ and a category k , $\frac{1}{2}x \oplus^k \frac{1}{2}y = z$ when there exists a, b such that $(x_i, a_{-i}) \sim^{k*} (z_i, b_{-i})$ and $(z_i, a_{-i}) \sim^{k*} (y_i, b_{-i})$. If \succsim^k has an additive representation, then $\frac{1}{2}U_i^k(x) + \frac{1}{2}U_i^k(y) = U_i^k(z)$. Define \oplus^k similarly for alternatives: $\frac{1}{2}x \oplus^k \frac{1}{2}y = z$ if and only if $z_i = \frac{1}{2}x_i \oplus_i^k \frac{1}{2}y_i$ for each dimension i . Finally, define $\alpha x \oplus^k (1 - \alpha)y$ by taking limits.¹⁰ We note that if U_i^k is linear, then $\alpha x \oplus_i^k (1 - \alpha)y = \alpha x + (1 - \alpha)y$.

Axiom 6 (Affine Across Categories (AAC)). For any $r \in X$, $x, x', \alpha x \oplus^j (1 - \alpha)x' \in K^j(r)$, and $y, y', \alpha y \oplus^k (1 - \alpha)y' \in K^k(r)$: if $x \succsim_r y$ and $x' \succsim_r y'$, then $\alpha x \oplus^j (1 - \alpha)x' \succsim_r \alpha y \oplus^k (1 - \alpha)y'$.

This axiom is a natural adaptation of the linearity axiom, a close relative of the independence axiom. If we strengthened Affine Across Categories to be stated using the traditional linearity condition, then we would obtain a representation where each $U^k(\cdot|r)$ is itself an affine function. Otherwise, it requires that the \oplus^k operation preserves indifference.

The second axiom deals with a technical issue.

Axiom 7 (Unbounded). For any $r \in X$: if $K^k(r)$ contains a sequence x_n so that $U^k(x_n) \rightarrow \infty$ ($-\infty$), then for any $x \in X$ there exists $x^* \in K^k(r)$ so that $x^* \succ_r x$ ($x \succ_r x^*$).

We note that U^k is unique up to a positive affine transformation. Hence whenever the utility of some sequence goes to infinity for some representation of \succsim^k , it must

¹⁰In general, $\alpha x \oplus^k (1 - \alpha)y$ need not exist. However, it does exist “locally,” which is all we require in the proof. That is, if $x \in K^k(r)$, then there exists an open set O with $x \in O$ on which $\alpha y \oplus^k (1 - \alpha)z$ exists for every $\alpha \in [0, 1]$ and $y, z \in O$.

also converge to infinity for any other representation as well. While the axiom can be stated in terms of primitives, we instead state it in terms of the U^k .¹¹ It ensures that a category containing alternatives whose utility goes to positive (negative) infinity must contain an alternative better (worse) than any other given alternative. If it failed, then no affine transformation of the category utility would represent the preference.

Theorem 3. *Assume the Structure Assumption holds. Then, $\{\succsim_r\}_{r \in X}$ satisfies the CTM axioms, Affine Across Categories, and Unbounded for \mathcal{K} if and only if it conforms to Affine Increasing CTM under \mathcal{K} .*

All the models discussed in Section 2 fall into the class of Affine CTM, so the result reveals the behavior all have in common. Relative to CTM, Affine Across Categories imposes stronger requirements on how the DM relates alternatives in different categories. Not only does the DM evaluate utility within a category using an additive function, but the additive structure persists across categories. Moreover, this aids with interpreting utility differences. If every pair of categories contains alternatives indifferent to one another, the entire representation is unique up to a common positive affine transformation. We call the combination of Axioms 1-4 and 6-7 the Affine CTM axioms.

3.3. Behavioral Foundation for Strong CTM. For a strong CTM, changing the reference point does not reverse the ranking of two products unless it also changes their categorization. The following axiom imposes this.

Axiom 8 (Reference Irrelevance). For any $x, y, r, r' \in X$:

if $x \in K^k(r) \cap K^k(r')$ and $y \in K^l(r) \cap K^l(r')$, then $x \succsim_r y$ if and only if $x \succsim_{r'} y$.

For the general CTM, the reference point influences choice through two channels: the category to which it belongs and its valuation. The axiom eliminates the latter.

¹¹ The statement in terms of primitives involves standard sequences and does not reveal key aspects of behavior, so we instead present the simpler and easier to interpret one above. In special cases, this is easy to do. For instance, if U^k is linear, then the axiom simply states that if $K^k(r)$ is an unbounded set, then the conclusion of the above axiom holds.

When comparing two alternatives across different reference points, the DM's relative ranking does not change when neither's category changes. This property greatly limits the effect of the reference point. In fact, a sufficiently small change in the reference never leads to a preference reversal.

Theorem 4. *Assume the Structure Assumption holds and for any categories i, j and any $r \in X$, there exists $x \in K^i(r)$ and $y \in K^j(r)$ with $x \sim_r y$. Then, $\{\succsim_r\}_{r \in X}$ satisfies the Affine CTM axioms and Reference Irrelevance for \mathcal{K} if and only if conforms to Strong, Increasing CTM under \mathcal{K} .*

Since BGS, MO, and PT are Strong CTM, Theorem 4 characterizes the behavior they have in common. While the reference plays a role in categorization, it plays no role in choice after categorization is taken into account. TK, which belongs to affine CTM but not strong CTM, must therefore violate reference irrelevance.

3.4. Comparing Models of Riskless Choice. TK, BGS, MO, PT, and the neo-classical model all conform to Affine CTM, so Theorems 1 and 3 describe the behavior that they have in common. However, the analysis so far, as well as the functional forms of the models, leaves open the question of what behavior distinguishes them. Of course, they differ in how alternatives are categorized, but the models also reflect distinct behavior within and across categories.

In addition to Reference Irrelevance, they are distinguished by whether they satisfy two classic axioms: Monotonicity and Cancellation, the unrestricted versions of Category Monotonicity and Category Cancellation.¹² The first requires that a dominant bundle is chosen, and the latter that an additive structure obtains. The representation theorem of Tversky & Kahneman [1991] imposes those two axioms in addition to continuity. In Appendix A.8, we show that an affine CTM with a Gain-Loss category function satisfies the two classic axioms and continuity if and only if it has a TK representation. We provide a detailed examination of the BGS model in Section 4.

¹²The formal statements are obtained by dropping the requirement in those two axioms that the alternatives belong to the same category.

Table 1 compares the four models in terms of Reference Irrelevance, Monotonicity and Cancellation, when BGS, TK, MO, and PT do not coincide with the neoclassical model. Only the neoclassical model satisfies all conditions; none of the other four do. On the one hand, BGS and PT satisfy Reference Irrelevance but violate Monotonicity and Cancellation. On the other, TK maintains Monotonicity and Cancellation but violates Reference Irrelevance. Finally, MO satisfies all but Cancellation.¹³

	Neoclassical	BGS	TK	MO	PT
CTM	✓	✓	✓	✓	✓
Monotonicity	✓	✗	✓	✓	✗
Reference Irrelevance	✓	✓	✗	✓ ¹⁴	✓
Cancellation	✓	✗	✓	✗	✗

TABLE 1. Comparisons of Models

We provide a plausible example violating the Cancellation axiom, and hence behavior inconsistent with TK. Then, we illustrate BGS can accommodate this example even without requiring a shift in the reference point. While the example is one simple test to distinguish BGS from TK, it is also powerful as it works for a fixed reference point.

Example 1. *Consider a consumer who visits the same wine bar regularly. The bartender occasionally offers promotions. The customer prefers to pay \$8 for a glass of French Syrah rather than \$2 for a glass of Australian Shiraz. At the same time, she prefers to pay \$2 for a bottle of water rather than \$10 for the glass of French Syrah. However, without any promotion in the store, she prefers paying \$10 for Australian Shiraz to paying \$8 for water.*

¹³Propositions 2 and 5 give the ✓'s of the table for BGS and TK. It is routine to verify that MO satisfies Monotonicity and Reference Irrelevance and the PT satisfies RI. We provide examples showing the other properties are violated in Appendix A.5.

¹⁴Whenever $c(r) = c(r')$ for every $r, r' \in X$.

The behavior in this example is both intuitively and formally consistent with the salient thinking model of BGS.¹⁵ Without any promotion, the consumer expects to pay a high price for a relatively low quality selection. When choosing between Syrah or Shiraz, the consumer focuses on the French wine's sublime quality, and she is willing to pay at least \$6 more for it. When choosing between water and Syrah, the low price of water stands out and she reveals that the gap between wine and water is less than \$8. However, when there is no promotion, she focuses again on the quality, and she is willing to pay an additional \$2 for even her less-preferred Australian Shiraz over water. Notice that this explanation does not require that the reference points are different. Since the consumer visits this bar regularly, intuitively, her reference point should be fixed and stable.

3.5. Non-increasing CTM. For simplicity, we have so far focused on increasing CTM. This is a desirable feature in consumer choice, but models of social preference often violate this property. For instance, inequality-averse individual 1 prefers to increase the allocation to individual 2 from x to y when she feels guilty but not when she is envious. However, she always prefers increasing the allocation to 2 in an allocation categorized as guilty, and to decrease in any categorized as envious. This contradicts Category Monotonicity, suggesting the following weakening.

Axiom (Consistent Preference within Category, CPC). *For each category k , there exists a set of attributes P^k so that if $x_j \geq y_j$ for all $j \in P^k$, $y_i \geq x_i$ for all $i \notin P^k$, and $x \neq y$, then $y \succ^{k*} x$.*

The set P^k contains the attributes for which an increase positively affects the DM's evaluation. CPC requires that the set of positive attributes in a category does not depend on the reference point. For the two-person-RIA model, the set for the

¹⁵Implicitly, the example reveals that the quality of French Syrah is higher than Australian Shiraz which is in turn higher than water. The numerical value of quality assigned to each beverage is irrelevant to the violation of Cancellation. For examples of qualities so that choice can be represented by the BGS model, one can calculate that $(-8, q_{fs}) \succ_r (-2, q_{as})$, $(-2, q_w) \succ_r (-10, q_{fs})$ and $(-10, q_{as}) \succ_r (-8, q_w)$ for $q_{fs} = 8$, $q_{as} = 6.9$, $q_w = 5.1$, and the reference point $r = (\frac{1}{2}(-10 + -8), \frac{1}{2}(q_w + q_{as}))$ when $w = 0.6$.

“guilty” category is $\{1, 2\}$ since she strictly prefers increasing everyone’s allocation, but the set for the “envious” one is $\{1\}$ – she prefers more for herself but dislikes others having even more. Note that CM is the special case of CPC where P^k includes every dimension for every category.

A CTM is characterized by all the properties of an increasing CTM, except where CM is replaced by CPC. The proof is a straightforward generalization of earlier one, so it is omitted.

4. BGS MODEL AND CATEGORIES

The BGS model is intuitive, tractable, and accounts for a number of empirical anomalies for the neoclassical model of choice. Despite its popularity, it can be difficult to understand all of the implications of the BGS model. Its new components are unobservable, and its functional form rather involved.

The first crucial step towards understanding the model is getting a handle on the novel salience function that determines which attribute stands out for a given reference point. While one can work out the implications of a particular salience function, this exercise is not fruitful since the particular function that applies to a given agent is unobservable. Moreover, it is not clear how the model changes when the underlying salience function changes.

CTM provides a lens through which we can study the salience function. While it influences which attribute is salient, the weight given to each attribute is independent of its magnitude. Therefore, its role is simply to divide the domain into distinct categories, each associated with a particular attribute being most salient. We study the salience function by focusing on the properties of the categories it generates.

Categories are generated by a function $s : \mathbb{R}_{++} \times \mathbb{R}_{++} \rightarrow \mathbb{R}_+$ if $x \in K^i(r)$ if and only if $s(x_i, r_i) > s(x_j, r_j)$ for all $j \neq i$. In the BGS model, categories are generated by a salience function σ that must satisfy the following properties. First, it increases in


contrast, i.e. for $\epsilon > 0$ and $a > b$, $\sigma(a + \epsilon, b) > \sigma(a, b)$ and $\sigma(a, b - \epsilon) > \sigma(a, b)$. Second, it is continuous in both arguments. Third, it is symmetric, i.e. $\sigma(a, b) = \sigma(b, a)$. Two other properties are sometimes assumed: σ is Homogeneous of Degree Zero (HOD) if for all $\alpha > 0$, $\sigma(\alpha a, \alpha b) = \sigma(a, b)$, and σ has diminishing sensitivity if for all $\epsilon > 0$ and $a, b > 0$, $\sigma(a + \epsilon, b + \epsilon) \leq \sigma(a, b)$.¹⁶ Finally, we always impose that the salience function is grounded: $\sigma(r, r) = \sigma(r', r')$ for all $r, r' \in X$. This is an implication of HOD satisfied by all of the specifications of which we are aware in the literature, and is a necessary condition for an attribute to be salient only if it differs from the reference.

Consider the following properties of categories.

- S0:** (Basic) For any $r \in X$: $K^1(r) \cap K^2(r) = \emptyset$, $K^1(r) \cup K^2(r)$ is dense in X , K^1, K^2 are continuous at r , and $K^1(r), K^2(r)$ are regular open sets.
- S1:** (Moderation) For any $\lambda \in [0, 1]$ and $r \in X$: if $x \in K^k(r)$, $y_k = x_k$, and $y_{-k} = \lambda x_{-k} + (1 - \lambda)r_{-k}$, then $y \in K^k(r)$.
- S2:** (Symmetry) If $(a, b) \in K^k(c, d)$, then $(c, d) \in K^k(a, b)$ and $(b, a) \in K^{-k}(d, c)$.
- S3:** (Transitivity) If $(a_1, a_2) \notin K^2(r_1, r_2)$ and $(a_2, a_3) \notin K^2(r_2, r_3)$ then $(a_1, a_3) \notin K^2(r_1, r_3)$.
- S4:** (Difference) For any x, y, z with $y \neq z$, $(x, y) \in K^2(x, z)$ and $(y, x) \in K^1(z, x)$.
- S5:** (Diminishing Sensitivity) For any $x, y, K^1, K^2, \epsilon > 0$, if $(x, y) \notin K^1(r_1, r_2)$, then $(x + \epsilon, y) \notin K^1(r_1 + \epsilon, r_2)$.
- S6:** (Equal Salience) For any $x, r \in X$: if $\frac{x_1}{r_1} = \frac{x_2}{r_2}$ or $\frac{x_1}{r_1} = \frac{r_2}{x_2}$, then $x \notin K^k(r)$ for $k = 1, 2$.

The properties have natural interpretations. Any category function satisfies **S0** by definition; we include it for completeness. **S1** indicates that making a bundle's less salient attribute closer to the reference point does not change the salience of the bundle. That is, when x and y differ only in attribute l , and y is closer to the reference in that attribute, if x is k -salient, then so is y . **S2** requires that the same ranking is used for

¹⁶BGS require this inequality to hold strictly. However, this is not a desirable property. If σ is HOD as they assume, then $\sigma(r, r) = \sigma(\alpha r, \alpha r) = \sigma(r + \epsilon, r + \epsilon)$ for $\alpha > 1$ and $\epsilon = (\alpha - 1)r$, violating their definition of diminishing sensitivity.

FIGURE 4. Properties **S0-S6** Illustrated

each attribute. **S3** adapts transitivity to the salience ranking. It says that if a_1 stands out more relative to r_1 than a_2 does to r_2 , and a_2 stands out more relative to r_2 than a_3 does to r_3 , then a_1 stands out more relative to r_1 than a_3 does to r_3 . **S4** says simply that any difference stands out more than no difference. **S5** implies that increasing both the good and the reference by the same amount in the same dimension does not move the good from one category to another. **S6** reads that if every attribute of x differs from the reference point by the same percentage, then none of the attributes stands out. More formally, if the percentage difference between x_k and r_k is the same across attributes, then x is not k -salient for any $k \in \{0, 1\}$.

Figure 4 provides examples satisfying some but not all of the properties. The functions that generate them, as well as a verification that they satisfy the claimed properties, can be found in Example 4 in the Appendix.

Theorem 5. *The category function satisfies:*

- (1) **S0-S4** if and only if there exists a salience function σ that generates them;
- (2) **S0-S5** if and only if the σ that generates it has diminishing sensitivity; and
- (3) **S0**, **S1**, and **S6** if and only if it satisfies **S0-S6** if and only if the σ that generates it is HOD. Any HOD salience function generates the same categories.

This theorem provides a characterization for BGS's salience function.¹⁷ It translates the functional form assumptions on the salience function in terms properties on

¹⁷Theorem 5 relies on the full structure of \mathbb{R}^2 for the last two results, as noted in Footnote 1. Diminishing sensitivity and Homogeneity are both cardinal properties, and so are undefined without

the salience categories. The most common specification of the salience function, HOD, satisfies all of the above properties. Surprisingly, the result shows that there is a unique category function satisfying these properties. Hence, any two HOD salience functions lead to exactly the same behavior.

We now turn to the question of identifying the salience function from choice behavior alone. That is, given that we observe a family $\{\succsim_r\}_{r \in X}$, can we identify which alternatives have what salience?

Proposition 1. *Suppose that $\{\succsim_r\}_{r \in X}$ has a BGS representation. Then, the weights, utility indices, and salience function are uniquely identified from $\{\succsim_r\}_{r \in X}$.*

The proof provides an algorithm for this in general. We illustrate for the case where u_1 and u_2 are linear. Fixing a reference point r , any alternative that differs only in dimension i from r must be i -salient. Hence, we can identify the weights on dimensions within each category from the slope of the indifference curve passing through that alternative. Now, we can test whether y is 1-salient by seeing if the indifference curves close to it are those generated by the weights for 1-salient alternatives. Varying y and r allows identification of the salience function, and hence the categories.

In addition to the particular form of categories, BGS satisfies several properties that distinguish it from other CTMs. The most general of these is Reference Irrelevance, above, making BGS a strong CTM. The other follows.

Axiom 9 (Salient Dimension Overweighted, SDO). For any $x, y, r, r' \in X$: if $x, y \in K^k(r) \cap K^l(r')$, $x \succsim_r y$, $x_l > y_l$, and $y_k > x_k$, then $x \succ_{r'} y$.

This axiom requires that categories correspond to the dimension that gets the most weight. That is, the DM is more willing to choose an alternative whose “best” attribute is k when it is k -salient. To illustrate, consider alternatives x, y with $x_1 > y_1$ and $y_2 > x_2$. Because x is relatively strong in attribute 1, x should benefit more than

cardinal structure on X . Properties **S0-S4** are defined. Subsequent results that rely on Theorem 5, such as Propositions 2 and 3, remain true when imposing only **S0-S4** in this setting.

y from a focus on it. If x is chosen over y when attribute 2 stands out for both, then this advantage in the first dimension is so strong that even a focus on the other one does not offset it. Hence, the DM should surely choose x over y for sure when attribute 1 stands out for it.

Proposition 2. *Assume that there exists $x \in K^k(r)$ and $y \in K^j(r)$ with $x \sim_r y$ for any categories k, j and any $r \in X$. Then, the family $\{\succsim_r\}_{r \in X}$ satisfies the Affine CTM axioms, Reference Interlocking, Reference Irrelevance, and Salient Dimension Overweighted for a category function \mathcal{K} satisfying **S0-S5** if and only if it has a BGS representation where σ has diminishing sensitivity.*

This result characterizes the BGS model. It also provides guidance for comparing it with other models in the CTM class (see Figure 1 and Table 1). By outlining the model's testable implications, the result provides guidance on how to design experiments to test it.¹⁸

In their 2013 paper, BGS focus on a special case where the model is linear: $w_1^1 = w_2^2 = 1 - w_1^2 = 1 - w_2^1 > \frac{1}{2}$ and $u_1(x) = u_2(x) = x$. In an earlier version of this paper, we show this model is characterized by strengthening Affine Across Categories to require linearity and imposing a reflection axiom that requires permuting two alternatives and the reference point in the same way not to reverse the DM's choice between the two.¹⁹

Taken together Propositions 1 and 2 provide an outline for a fully subjective axiomatization of a family of preferences with a BGS representation. Proposition 1 shows that we can reveal a category function from the family of preferences, provided they have a representation. We check whether these revealed categories exist and satisfy S0-S5. If so, then the axioms shown necessary by the second result apply with this revealed category function.

¹⁸The assumption that alternatives indifferent to each other exist in each category for each reference point is not strictly necessary. A sufficient condition for it to be necessary is that the utility indexes are both unbounded above (or below).

¹⁹Formally, the first is that Affine Across Categories holds with \oplus^k replaced by the usual $+$ operation. The second is that $(a, b) \succsim_{r_1, r_2} (c, d)$ if and only if $(b, a) \succsim_{r_2, r_1} (d, c)$. One can verify that these additional assumptions imply that the ancillary assumption about indifference holds.

5. CHOICE CORRESPONDENCE

In this section, the modeler observes only the DM’s choice from a finite subset of choices and nothing more. A model consists of both a theory of reference formation and a theory of choice given categorization. In this setting, we can jointly test the theory of choice given categorization, categorization given reference, and reference formation.

We model reference formation via a reference generator, a map from finite subsets of alternatives to reference points. We denote the reference generator $A : 2^S \setminus \emptyset \rightarrow X$, with the interpretation that $A(S)$ is the reference point when the menu is S . Examples include the BGS theory that $A(S)$ is the average alternative, that $A(S)$ is the median bundle, that $A(S)$ is the upper (or lower) bound of S , and the Köszegi & Rabin [2006] theory that $A(S) = c(S)$. If additional observable data on the choice context is provided, then it is easy to extend our results to A being a function of that as well. For instance MO theorize that the initial endowment e is observable and that $A(S, e) = e$, and Bordalo et al. [2019] theorize that past histories h of consumption are available and that $A(S, h)$ is the average between the bundles in S and those in h .

Fixing a categorization function \mathcal{K} and a reference generator A , let \mathcal{X} be the set of finite and non-empty subsets of X such that every alternative is categorized. Formally, $S \in \mathcal{X}$ only if $S \subset \bigcup_{i=1}^m K^i(A(S))$. We call these menus or categorized menus for short. The requirement ensures that each alternative in the choice set belongs to a category given the reference point $A(S)$. We leave open how the DM chooses when alternatives that are uncategorized belong to the choice set. By leaving the choice from this small set of menus ambiguous, we can more clearly state the properties of choice implied by the model.²⁰

²⁰One can, of course, extend the model to account for these choices. For instance, BGS hypothesize that these alternatives are evaluated according to their sum. Complications arise because the uncategorized alternatives are “small:” its complement is open and dense, and moreover it has zero measure.

We summarize the DM's choices by a choice correspondence $c : \mathcal{X} \rightrightarrows X$ with $c(S) \subseteq S$ and $c(S) \neq \emptyset$ for each $S \in \mathcal{X}$. Adapted to this setting, the model has the following representation.

Definition 4. The choice correspondence c *conforms to Strong-CTM under* (\mathcal{K}, A) if there exists a family of preference relations $\{\succsim_r\}_{r \in X}$ that conforms to Increasing Strong CTM under \mathcal{K} so that

$$c(S) = \left\{ x \in S : x \succsim_{A(S)} y \text{ for all } y \in S \right\}$$

for every $S \in \mathcal{X}$.

5.1. Reference point formation. Provided that the reference generator is responsive enough to changes in the menu, there is the possibility of testing the properties required by categorization on \succsim_r . One example of enough structure is that the reference point is the average bundle. However, this is just one example. An even more general sufficient condition is as follows.

Assumption. A function A is a *generalized average* if for any $S = \{x^1, \dots, x^m\} \in \mathcal{X}$:

- (i) the function $x \mapsto A([S \setminus \{x_1\}] \cup \{x\})$ is continuous at x_1 , and
- (ii) for any $\epsilon > 0$ and any finite $S' \in \bigcup_i K^i(A(S))$, there exists $S^* \in \mathcal{X}$ so that $S^* \supset S \cup S'$, $d(A(S^*), A(S)) < \epsilon$, and for any $x' \in S^* \setminus S'$, $\min_{x \in S} d(x', x) < \epsilon^2$.

Examples of generalized average reference include the average bundle

$$A_a(S) = \left(\frac{\sum_{x \in S} x_1}{|S|}, \frac{\sum_{x \in S} x_2}{|S|} \right),$$

the median value of each attribute, and a weighted average

$$A_{wa}(S) = \left(\frac{\sum_{x \in S} w(x)x_1}{\sum_{x \in S} w(x)}, \frac{\sum_{x \in S} w(x)x_2}{\sum_{x \in S} w(x)} \right)$$

for any continuous weight function $w : X \rightarrow [a, b]$ with $b > a > 0$. We sometimes impose the additional requirement that $A(S) \in co(S) \setminus ext(S)$ for all non-singleton S ; if so, we call A a *strong generalized average*. The first and last of these examples satisfy

this property. The supremum and infimum on their own are *not* weighted averages, nor (necessarily) is the choice acclimating reference generator, $c(S) = A(S)$.²¹

5.2. Behavioral Foundations for Strong-CTM. We now consider the behavior by a DM who conforms to Strong-CTM for a given category function and reference generator. To do so, we make use of our earlier analysis by revealing how the DM evaluates alternatives categorized in a given way. When $A(S)$ is a generalized average, this provides enough structure to identify enough of the family to apply our earlier analysis.

The main behavioral content comes from the choice correspondence equivalent of Reference Irrelevance. To state it, we introduce the following definition and notation.

Definition 5. The alternative x in category k is indirectly revealed preferred to alternative y in category j , written $(x, k) \succsim^R (y, j)$, if there exists finite sequences of pairs $(x^i, S^i)_{i=1}^n$ such that $x = x^1 \in K^k(A(S^1))$, $y \in K^j(A(S^n)) \cap S^n$, and for each i : $x^i \in c(S^i)$, $x^{i+1} \in S^i$, and $x^{i+1} \in K^{k_i}(A(S^i)) \cap K^{k_i}(A(S^{i+1}))$ for some k_i .

We replace Reference Irrelevance with the following weakening of the Strong Axiom of Revealed Preference (SARP).

Axiom (Category SARP). For any $S \in \mathcal{X}$, if $(x, k) \succsim^R (y, j)$, $x \in K^k(A(S)) \cap S$, $y \in K^j(A(S)) \cap S$, and $y \in c(S)$, then $x \in c(S)$.

We first illustrate in a simple two menu setting, analogous to a test case for the Weak Axiom of Revealed Preference (WARP). Consider two menus S^1 and S^2 and two chosen products $x^1 \in c(S^1)$ and $x^2 \in c(S^2)$ where both products are categorized in the same way for both menus. For example, x^1 is in category 1 for both menus, and x^2 is in category 2 for both. The observation $x^1 \in c(S^1)$ reveals that the valuation of x^1 is at least as high as that of x^2 when x^1 belongs to the first category and x^2 to the second. Since the categorization of products does not change when the menu changes

²¹Recall $\sup S = (\max_{x \in S} x_1, \max_{x \in S} x_2)$ and $\inf S$ is defined analogously.

from S^1 to S^2 , their relative valuation stays the same as well. Hence, if x^2 is chosen from S^2 , then x^1 must be chosen too. Since neither products' category has changed, the DM should obey WARP for these two menus. However, the axiom leaves open the possibility of a WARP violation when either is differentially categorized.

The axiom extends this logic to sequences of choices in much the same way that SARP does to WARP. A finite sequence of choices, where the choice from the next menu is available in the current one and has the same salience in both, does not lead to a choice reversal. Since salience does not change along the sequence of choices, the choices do not exhibit a reversal.

Category SARP limits the effect of unchosen alternatives. Modifying them can alter the DM's choice, but only insofar as changing them changes the reference point and thus the salience of alternatives. It states that these unchosen options do not alter the relative ranking of two alternatives, unless they change the region to which the alternatives belong. That is, when comparing the same two alternatives in different menus, the DM's relative ranking does not change when neither's salience changes. This property greatly limits the effect of the reference point. In fact, a sufficiently small change in the reference never leads to a preference reversal.

The remaining axioms are the natural generalizations to the choice correspondence of Category Cancellation, Category Monotonicity, Category Continuity, Reference Interlocking, and Affine Across Categories. We denote these by appending a “*” to distinguish from their reference-dependent-preference formulation. Appendix B.1 contains their formal statement.

As before, we require some additional topological structure on the categories. For a category k , let

$$E^{R,k} = \{x \in X : x \in K^k(A(S)), \{x\} = c(S)\}$$

and

$$D^k = \bigcup_{S \in \mathcal{X}} \{K^k(A(S)) \cap S\}.$$

The generalization of the structure assumption is as follows.

Assumption (Revealed Structure). For any category k , $E^{R,k}$ is open, $E^{R,k}$ is dense in D^k , and the following sets are connected: $E^{R,k}$, $\{x \in E^{R,k} : x_j = s\}$ for all dimensions j and scalars $s \in \mathbb{R}$, and $\{y \in E^{R,k} : (x, k) \sim^R (y, k)\}$ for all $x \in E^{R,k}$.

In addition to what was imposed by the Structure Assumption, we require that almost all objects categorized in a category are chosen in some menu. This can be weakened, but is typically satisfied by the models in which we are interested, such as BGS.

We require one last assumption.

Axiom (Comparability Across Regions, CAR). If $x \in E^{R,k}$, then for any j there exists $y \in E^{R,j}$ so that $(x, k) \sim^R (y, j)$.

This is a version of the assumption we made for Strong CTM. It requires that every alternative chosen when it belongs to category k is revealed to be equally good to some other alternative when it is categorized in category j . With it, we can now state the result.

Theorem 6. Assume that Revealed Structure and CAR hold and that A is a generalized average. A choice correspondence c conforms to strong-CTM under (\mathcal{K}, A) if and only if c satisfies Category-SARP, Category Monotonicity*, Category Cancellation*, Category Continuity*, and Affine Across Categories*.

The result is the counterpart of Theorem 4 with an endogenous reference point. The behavior corresponding to categorization does not fundamentally change across settings. As long as the DM reacts consistently when alternatives are categorized in the same way, then we can represent her choices as categorical thinking where the

reference point only affects how she categorizes each alternative. The key challenge in the proof is to establish that the arguments we used to establish our earlier results still hold. We adapt our earlier arguments to show that revealed preference within category k is complete on $E^{R,k}$. This relies on small changes in alternatives not changing choice, a property implied by generalized average. Then, the remaining axioms establish that this within-category preference has an additive representation. CAR allows us to extend across categories.

5.3. Behavioral Foundations for BGS. In this subsection, we provide a behavioral foundation for BGS. The first step is to show that the Revealed Structure assumption holds.

Lemma 1. *If A is a strong generalized average, \mathcal{K} satisfies **S0**, **S1**, and **S4**, and c satisfies Category Monotonicity*, then $E^{R,k} = \mathbb{R}_{++}^2$ for $k = 1, 2$.*

Given the assumptions we have made so far, every alternative is chosen in some menu when it is k -salient. Consequently, the revealed structure assumption must hold. The result relies on the observation that the DM categorizes x as 1-salient when all other available options have the same value in dimension 2 as x . If x has the highest value in attribute 1 in such a choice set, then it must be chosen.

Now, we can apply Theorem 6 in combination with the insights gained from Proposition 2 to understand the behavioral foundation of the BGS model.

Proposition 3. *Assume that A is a strong generalized average and that CAR holds. The choice correspondence c satisfies Category-SARP, Category Monotonicity*, Category Cancellation*, Category Continuity*, Affine Across Categories*, Reference Interlocking*, and Salient Dimension Overweighted* for a category function \mathcal{K} satisfying **S0-S5** if and only if c conforms to BGS where σ has diminishing sensitivity.*

This theorem lays out the behavioral postulates that characterize the BGS model with endogenous reference point formation. Most importantly, it connects the (unobserved) components of the model to observed choice behavior. Fundamentally, the properties that Proposition 2 characterized the model in our first setting still characterize it. To do so, we note that Theorems 5 and 6 imply that there exists a Strong CTM with categories generated by a salience function. We then establish that choice within the k -salient alternatives overweights dimension k by using SDO and the structure of regions.

Finally, we ask the question of whether the choice correspondence with an endogenous reference point provides enough leverage to identify salience.

Proposition 4. *Given that c conforms to BGS with a strong generalized average, the categories are uniquely identified.*

As with Propositions 1 and 2, Propositions 3 and 4 provide a roadmap for testing BGS without a known salience function. However, it still requires that the reference generator is a strong generalized average. Consequently, the axioms capture the full testable implication of the model and allow for tight comparisons with other existing work.

6. RELATED LITERATURE

This paper provides a choice theoretic analysis of categorization. We apply this model to highlight similarities and differences between a number of behavioral models in the literature. As such, it is closely related to the literature which studies how a reference point affects choices, (*e.g.* Tversky & Kahneman [1991], Munro & Sugden [2003], Sugden [2003], Masatlioglu & Ok [2005], Sagi [2006], Salant & Rubinstein [2008], Apesteguia & Ballester [2009], Masatlioglu & Nakajima [2013], Masatlioglu & Ok [2014], Dean, Kıbrıs, & Masatlioglu [2017]). The papers focus on an exogenous

reference point, as in Section 3. While TK and MO are examples of CTM, the others are not. Nonetheless, our analysis puts the models on an equal footing so their implications can be compared.

We then extend the model to consider endogenous reference point formation. This adopts the approach of a number of recent papers, *e.g.* Bodner & Prelec [1994], Kivetz, Netzer, & Srinivasan [2004], Orhun [2009], Bordalo, Gennaioli, & Shleifer [2012], Tserenjigmid [2015]. As in Section 5, the reference point is a function of the context, and is identical for all feasible alternatives. Finally, Köszegi & Rabin [2006], Ok, Ortoleva, & Riella [2015], Freeman [2017] and Kıbrıs et al. [2018] study models where the endogenous reference point is determined by what the agent chooses, but is otherwise independent of the choice set. This represents a very different approach to reference formation, and our approach does not easily generalize to accommodate it.²²

One of our key contributions is to provide an axiomatization of the salient thinking model. Interpreting salience as arising from differential attention to attributes, CTM has a close relationship with the literature studying how limited attention affects decision making. Masatlioglu et al. [2012] and Manzini & Mariotti [2014] study a DM who has limited attention to the alternatives available. The DM maximizes a fixed preference relation over the consideration set, a subset of the alternatives actually available. In contrast, in CTM the DM considers all available alternatives but maximizes a preference relation distorted by her attention. Caplin & Dean [2015], de Oliveira et al. [2017] and Ellis [2018] study a DM who has limited attention to information. In contrast to CTM, attention is chosen rationally to maximize ex ante utility, rather than determined by the framing of the decision, and choice varies across states of the world. The most related interpretation considers attributes as payoffs in a fixed state. In addition to choices varying across states, each alternative has the same weights on each attribute, similar to Köszegi & Szeidl [2013]. Taken together, these results highlight the effects on behavior of different types of attention.

²²Maltz [2017] is the only model of which we are aware that combines an exogenous reference point with endogenous reference-point formation.

While we argue in this paper that a number of prominent behavioral economic models can be thought of as resulting from categorization, few papers in economics explicitly address categorization. Mullainathan [2002] provides a model of belief updating and shows how categorization can generate non-Bayesian effects. Fryer & Jackson [2008] introduce a categorical model of cognition where a decision maker categorizes her past experiences. Since the number of categories is limited, the decision maker must group distinct experiences in the same category. In this model, prediction is based on the prototype from the category which matches closely the current situation. Finally, Manzini & Mariotti [2012] introduce a two-stage decision-making model. In the first stage, a decision maker eliminates some of alternatives based on their categories, and in the second stage she maximizes her preference among the alternatives surviving after the first stage. Bordalo et al. [2019] provide a model of memory and attention, where the context’s similarity to past consumption opportunities affects the salience of the alternatives currently available. They show this leads to endogenous categorization of the current opportunity set, and discuss the resulting implications for choice.

The evolutionary psychology literature on categorization suggests a common explanation for the effects shown to be captured by our model of categorization. That literature stresses that categories evolved as cues to apply a particular mental process in a given situation (see e.g. the review by Cosmides & Tooby [2013]). However, these processes are often applied to situations different from their evolutionary purpose. Boyer & Barrett [2015] explain, “*The fact that some cognitive system is specialized for a domain D does not entail that it invariably or exclusively handles D, nor does it mean that the specialization cannot be co-opted for evolutionarily novel activities.*” This implies that systems used to evaluate categorized objects are miscalibrated from how they would be more useful. For instance, New et al. [2007] documented that subjects were quicker and more accurate in noticing changes involving animals than for those involving vehicles, despite the latter’s much greater importance in modern life.

REFERENCES

- Anderson, J. R. (1991). The adaptive nature of human categorization. *Psychological review*, 98(3), 409.
- Apesteguia, J., & Ballester, M. A. (2009). A theory of reference-dependent behavior. *Economic Theory*, 40(3), 427–455.
- Ashby, F., & Gott, R. (1988). Decision rules in the perception and categorization of multidimensional stimuli. *Journal of experimental psychology. Learning, memory, and cognition*, 14, 33–53.
- Ashby, F. G., & Maddox, W. T. (2005). Human category learning. *Annual Review Psychology*, 56, 149–178.
- Ashby, F. G., & Waldron, E. M. (1999). On the nature of implicit categorization. *Psychonomic Bulletin and Review*, 6, 363–378.
- Bhatia, S., & Golman, R. (2013). Attention and reference dependence. *Working paper*.
- Bodner, R., & Prelec, D. (1994). The centroid model of context dependent choice. *Unpublished Manuscript, MIT*.
- Bordalo, P., Gennaioli, N., & Shleifer, A. (2012). Salience theory of choice under risk. *The Quarterly Journal of Economics*, 127(3), 1243–1285.
- Bordalo, P., Gennaioli, N., & Shleifer, A. (2013). Salience and consumer choice. *Journal of Political Economy*, 121(5), 803–843.
- Bordalo, P., Gennaioli, N., & Shleifer, A. (2019). Memory, attention, and choice. *Working Paper, Harvard*.
- Boyer, P., & Barrett, H. C. (2015). Intuitive ontologies and domain specificity. *The handbook of evolutionary psychology*, (pp. 1–19).
- Bushong, B., Rabin, M., & Schwartzstein, J. (2015). A model of relative thinking. *Unpublished manuscript, Harvard University, Cambridge, MA*.
- Caplin, A., & Dean, M. (2015). Revealed preference, rational inattention, and costly information acquisition. *American Economic Review*, 105(7), 2183–2203.
- Charness, G., & Rabin, M. (2002). Understanding social preferences with simple tests. *The Quarterly Journal of Economics*, 117(3), 817–869.

- Chateauneuf, A., & Wakker, P. (1993). From local to global additive representation. *Journal of Mathematical Economics*, 22(6), 523–545.
- Chernev, A. (2011). The dieter’s paradox. *Journal of Consumer Psychology*, 21(2), 178–183.
- Choi, J., & Kim, S. (2016). Is the smartwatch an it product or a fashion product? a study on factors affecting the intention to use smartwatches. *Computers in Human Behavior*, 63, 777 – 786.
- Cosmides, L., & Tooby, J. (2013). Evolutionary psychology: New perspectives on cognition and motivation. *Annual review of psychology*, 64, 201–229.
- de Oliveira, H., Denti, T., Mihm, M., & Ozbek, K. (2017). Rationally inattentive preferences and hidden information costs. *Theoretical Economics*, 12(2), 621–654.
- Dean, M., Kibris, Ö., & Masatlioglu, Y. (2017). Limited attention and status quo bias. *Journal of Economic Theory*, 169, 93–127.
- Debreu, G. (1959). Topological methods in cardinal utility theory. Tech. rep., Cowles Foundation for Research in Economics, Yale University.
- Ellis, A. (2018). Foundations for optimal inattention. *Journal of Economic Theory*, 173, 56–94.
- Fehr, E., & Schmidt, K. M. (1999). A theory of fairness, competition, and cooperation. *Quarterly Journal of Economics*, 114(3), 817–868.
- Freeman, D. J. (2017). Preferred personal equilibrium and simple choices. *Journal of Economic Behavior & Organization*, 143, 165–172.
- Fryer, R., & Jackson, M. O. (2008). A categorical model of cognition and biased decision making. *The BE Journal of Theoretical Economics*, 8(1).
- Gabaix, X. (2014). A sparsity-based model of bounded rationality, applied to basic consumer and equilibrium theory. *Quarterly Journal of Economics*, 130, 1369–1420.
- Ghirardato, P., Maccheroni, F., Marinacci, M., & Siniscalchi, M. (2003). A subjective spin on roulette wheels. *Econometrica*, 71(6), 1897–1908.
- Kibris, Ö., Masatlioglu, Y., & Suleymanov, E. (2018). A theory of reference point formation. Working papers.

- Kivetz, R., Netzer, O., & Srinivasan, V. (2004). Alternative models for capturing the compromise effect. *Journal of Marketing Research*, 41(3), 237–257.
- Kőszegi, B., & Rabin, M. (2006). A model of reference-dependent preferences. *Quarterly Journal of Economics*, 121(4), 1133–1165.
- Kőszegi, B., & Szeidl, A. (2013). A model of focusing in economic choice. *The Quarterly Journal of Economics*, 128(1), 53–104.
- Krantz, D., Luce, D., Suppes, P., & Tversky, A. (1971). *Foundations of measurement, Vol. I: Additive and polynomial representations*. New York Academic Press.
- Loken, B. (2006). Consumer psychology: categorization, inferences, affect, and persuasion. *Annu. Rev. Psychol.*, 57, 453–485.
- Loken, B., Barsalou, L. W., & Joiner, C. (2008). Categorization theory and research in consumer psychology. *Handbook of consumer psychology*, (pp. 133–165).
- Love, B. C., Medin, D. L., & Gureckis, T. M. (2004). Sustain: a network model of category learning. *Psychological review*, 111(2), 309.
- Maltz, A. (2017). Exogenous endowment - endogenous reference point. Working Papers WP2016/5, University of Haifa, Department of Economics.
- Manzini, P., & Mariotti, M. (2012). Categorize then choose: Boundedly rational choice and welfare. *Journal of the European Economic Association*, 10(5), 1141–1165.
- Manzini, P., & Mariotti, M. (2014). Stochastic choice and consideration sets. *Econometrica*, 82(3), 1153–1176.
- Masatlioglu, Y., & Nakajima, D. (2013). Choice by iterative search. *Theoretical Economics*, 8(3), 701–728.
- Masatlioglu, Y., Nakajima, D., & Ozbay, E. Y. (2012). Revealed attention. *American Economic Review*, 102(5), 2183–2205.
- Masatlioglu, Y., & Ok, E. A. (2005). Rational choice with status quo bias. *Journal of Economic Theory*, 121, 1–29.
- Masatlioglu, Y., & Ok, E. A. (2014). A canonical model of choice with initial endowments. *The Review of Economic Studies*, 81(2), 851–883.
- Mogilner, C., Rudnick, T., & Iyengar, S. S. (2008). The mere categorization effect: How the presence of categories increases choosers’ perceptions of assortment variety

- and outcome satisfaction. *Journal of Consumer Research*, 35(2), 202–215.
- Mullainathan, S. (2002). Thinking through categories. *NBER working paper*.
- Munro, A., & Sugden, R. (2003). On the theory of reference-dependent preferences. *Journal of Economic Behavior & Organization*, 50(4), 407–428.
- New, J., Cosmides, L., & Tooby, J. (2007). Category-specific attention for animals reflects ancestral priorities, not expertise. *Proceedings of the National Academy of Sciences*, 104(42), 16598–16603.
- Ok, E. A., Ortoleva, P., & Riella, G. (2015). Revealed (p)reference theory. *American Economic Review*, 105(1), 299–321.
- Orhun, A. Y. (2009). Optimal product line design when consumers exhibit choice set-dependent preferences. *Marketing Science*, 28(5), 868–886.
- Posner, M. I., & Keele, S. W. (1970). Retention of abstract ideas. *Journal of Experimental psychology*, 83(2p1), 304.
- Ratneshwar, S., & Shocker, A. D. (1991). Substitution in use and the role of usage context in product category structures. *Journal of Marketing Research*, 28(3), 281–295.
- Rosch, E. (1975). Cognitive representations of semantic categories. *Journal of Experimental Psychology: General*, 104(3), 192–233.
- Sagi, J. S. (2006). Anchored preference relations. *Journal of Economic Theory*, 130(1), 283–295.
- Salant, Y., & Rubinstein, A. (2008). (a, f): Choice with frames. *The Review of Economic Studies*, 75(4), 1287–1296.
- Savage, L. J. (1954). *The Foundations of Statistics*. Dover.
- Stewart, N., Brown, G. D., & Chater, N. (2002). Sequence effects in categorization of simple perceptual stimuli. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 28(1), 3.
- Strotz, R. (1955). Myopia and inconsistency in dynamic utility maximization. *The Review of Economic Studies*, 23(3), 165–180.
- Sugden, R. (2003). Reference-dependent subjective expected utility. *Journal of Economic Theory*, 111(2), 172–191.

- Tserenjigmid, G. (2015). Choosing with the worst in mind: A reference-dependent model. Working papers.
- Tversky, A. (1977). Features of similarity. *Psychological review*, 84(4), 327.
- Tversky, A., & Gati, I. (1978). Studies of similarity. *Cognition and categorization*, 1(1978), 79–98.
- Tversky, A., & Kahneman, D. (1991). Loss aversion in riskless choice: A reference-dependent model. *The Quarterly Journal of Economics*, 106(4), 1039–1061.
- Wanke, M., Bless, H., & Schwarz, N. (1999). Lobster, wine and cigarettes: Ad hoc categorisations and the emergence of context effects. *Marketing Bulletin*, 10, 52–56.

APPENDIX A. PROOFS AND EXTRAS FROM SECTIONS 2 - 4

A.1. Proof of Theorem 1.

Lemma 2. \succ^{k*} has open upper and lower contour sets in E^k .

Proof. Suppose $x \succ^{k*} y$. Then, there are $x^1, x^2, \dots, x^M \in E^k$ and r^1, \dots, r^M with $x^1 = x$ and $x^M = y$ so that $x^j \succ_{r^j} x^{j+1}$ and $x^j, x^{j+1} \in K^k(r^j)$. Let $\epsilon_j > 0$ be such that $B_{\epsilon_j}(x^j), B_{\epsilon_j}(x^{j+1}) \subset K^k(r^j)$. Set $\epsilon = \min\{\epsilon_j\}_{j < M}$.

Now, $x^j \succ^k x^{j+1}$ for at least one j . Let m be the last index for which this is true. Since $B_\epsilon(x^m), B_\epsilon(x^{m+1}) \subset K^k(r^m)$, Let $0 < \epsilon_m^* < \epsilon$ be such that $B_{2\epsilon_m^*}(x^m)$ is a subset of $\{x : x \succ_{r^m} x^{m+1}\}$. Then, $x^m - \epsilon_m^* \succ^k x^{m+1}$ and by definition of \succ^{k*} , it follows that $x^m - \epsilon_m^* \succ^{k*} y$. Assume (IH) that there is $0 < \epsilon_{m-j}^* < \epsilon$ so that $x^{m-j} - \epsilon_{m-j}^* \succ^{k*} y$. Then, it is easy to see that

$$x^{m-j-1} \succ^k x^{m-j} - \epsilon_{m-j}^*$$

for $j \geq 0$ using CM, weak order, and the construction of ϵ_{m-j}^* . By Category Continuity and Monotonicity, there then exists $0 < \epsilon_{m-j-1}^* < \epsilon$ so that $x^{m-j-1} - \epsilon_{m-j-1}^* \succ^k x^{m-j} - \epsilon_{m-j}^*$. By (IH) and the definition of \succ^{k*} , it follows that $x^{m-j-1} - \epsilon_{m-j-1}^* \succ^{k*} y$. By induction, there is ϵ_1^* so that $x^1 - \epsilon_1^* \succ^{k*} y$, so by CM and definition of \succ^{k*} , we have $x' \succ^{k*} y$ for any $x' \in B_{\epsilon_1^*}(x)$. Conclude the upper-contour set is open; similar arguments work for the lower-contour set. \square

Lemma 3. \succ^{k*} is complete on E^k .

Proof. Pick any $x, y \in E^k$ and let $E^* = E^k \cap B_{d(x,y)+1}(x)$. As the intersection of two intersecting connected sets, E^* is connected, and as a subset of \mathbb{R}^n , there is a continuous path $\theta : [0, 1] \rightarrow E^*$ so that $\theta(0) = x$ and $\theta(1) = y$. This θ can be chosen so that it crosses each \succ^{k*} indifference curve at most once. To see why, suppose that $\theta(a) \sim^{k*} \theta(b)$ and $b > a$. Since $IC = \{b' \in E^* : b' \sim^{k*} \theta(a)\}$ is path-connected, there is another continuous path $\theta' : [0, 1] \rightarrow IC$ with $\theta'(0) = \theta(a)$ and $\theta'(1) = \theta(b)$. Then the path θ^* given by $\theta^*(x) = \theta(x)$ for $x \notin [a, b]$ and $\theta^*(x) = \theta'\left(\frac{x-a}{b-a}\right)$ for $x \in [a, b]$ is also a continuous path from x to y . Constructing this for $a^* = \min\{a' : \theta(a') \sim^{k*} \theta(a)\}$ and $b^* = \max\{a' : \theta(a') \sim^{k*} \theta(a)\}$ gives a path that crosses IC at most once. These are well-defined since θ is continuous.

Now, let $Y = \theta^{-1}([0, 1])$. Y is closed since θ is continuous and so compact as a subset of $cl(B_{d(x,y)+1}(x))$. For any $z \in Y$, there exists $r_z \in X$ and $\epsilon_z > 0$ so that $B_z = B_{\epsilon_z}(z) \subset K^k(r_z)$. Since $B_z \subset K^k(r_z)$ and \succ^k is a subrelation of \succ^{k*} , \succ^{k*} is complete and transitive when restricted to B_z . Then, the collection $\{B_z : z \in Y\}$ is an open cover of Y and hence has a finite subcover $B_{z_1}, B_{z_2}, \dots, B_{z_m}$. W.L.O.G., B_{z_j}

is not a subset of B_{z_j} , for any j, j' and $\theta(z_j) < \theta(z_{j+1})$, so $x \in B_{z_1}$ and $y \in B_{z_m}$. Moreover, since θ crosses each indifference curve only once, if $z_k \succ^{k*} z_{k+1}$ ($z_k \prec^{k*} z_{k+1}$) for any k , then $z_j \succ^{k*} z_{j'}$ ($z_k \prec^{k*} z_{k+1}$) for any $j' > j$. W.L.O.G. consider the former. Pick $a^1 \in B_{z_1} \cap B_{z_2} \cap Y$ so that $x \succ^k a^1$ and then pick $a^j \in B_{z_j} \cap B_{z_{j+1}} \cap Y$ so that $a^{j-1} \succ^k a^j$. Then,

$$x \succ^{k*} a^1 \succ^{k*} a^2 \succ^* \dots \succ^{k*} a^m \succ^{k*} y.$$

Since \succ^{k*} is transitive, we conclude $x \succ^{k*} y$. Since x, y were arbitrary, \succ^{k*} is complete. \square

Apply CW Theorem 2.2 to get an additive representation $U^i(x)$ on E^i . For any $x, y \in K^i(r)$, $x \succsim_r y$ if and only if $U^i(x) \geq U^i(y)$ and $U^i(x) = \sum_j U_j^i(x_j)$.

Lemma 4. *For categories $K^i(r)$ and $K^j(r)$, either (i) there exists $x^i \in K^i(r)$ and $x^j \in K^j(r)$ so that $x^i \sim_r x^j$; or (ii) $x^i \succ_r x^j$ for all $x^i \in K^i(r)$ and $x^j \in K^j(r)$; or (iii) $x^j \succ_r x^i$ for all $x^i \in K^i(r)$ and $x^j \in K^j(r)$.*

Proof. If neither (ii) nor (iii) holds, then after relabeling categories if necessary, there exist $x \in K^i(r)$ and $y, z \in K^j(r)$ such that $y \succ_r x \succ_r z$. Let $UC_j(x)$ and $LC_j(x)$ be the strict upper and lower contour sets of x in category j for reference r . Any point in $K^j(r) \setminus [UC_j(x) \cup LC_j(x)]$ is indifferent to x , so either (i) holds or the set is empty. There exists an $\epsilon > 0$ such that for every $x' \in B_\epsilon(x)$, $y \succ_r x' \succ_r z$ by Category Continuity and hence $K^j(r) \neq U_j(x')$ and $K^j(r) \neq L_j(x')$. By Category Continuity, there exists $x' \in B_\epsilon(x)$ such that $K^j(r) \setminus [UC_j(x') \cup LC_j(x')] \neq \emptyset$ (otherwise, $B_\epsilon(x)$ is contained in the interior of the set considered), so we can take $y' \in K^j(r) \setminus [UC_j(x') \cup LC_j(x')]$ and conclude $y' \sim_r x'$. \square

Definition 6. A finite sequence (Q_1, \dots, Q_{m+1}) with each $Q_i \in \{K^1(r), \dots, K^n(r)\}$ is an *indifference sequence for r* (IS) if there exists $x^1, \dots, x^m, y^1, \dots, y^m$ with $x^k \in Q_k$, $y^k \in Q_{k+1}$ and $x^k \sim_r y^k$.

We omit the dependence on r when clear from context.

Define the relation \bowtie_r by $x \bowtie_r y$ if there exists an indifference sequence of categories (Q_1, \dots, Q_m) with $x \in Q_1$ and $y \in Q_m$. It is easy to see that \bowtie_r is an equivalence relation (reflexive, symmetric, and transitive). Let $[x]_r$ denote the \bowtie_r equivalence class of x .

Lemma 5. *If $y \notin [x]_r$ and $x \succ_r y$, then $x' \succ_r y'$ for all $x' \in [x]_r$ and $y' \in [y]_r$.*

Proof. Fix $x, y, r \in X$ with $y \notin [x]_r$ and $x \succ_r y$, and assume $x \in K^k$. Pick any $y' \in [y]_r$. By definition, there is an IS (Q_1, \dots, Q_m) with $y' \in Q_m$ and $y \in Q_1$. Let $i = 1$ and $y_1 = y$. If there exists $y'' \in Q_i$ with $y'' \succsim_r x$, then $y'' \succsim_r x \succ_r y_i$, so by Lemma 4,

we can find $z \in Q_i$ and $x' \in K^k$ with $z \sim_r x'$. If that occurs, then (K^k, Q_i, \dots, Q_1) is an IS and $y \in [x]_r$, a contradiction. Thus $x \succ_r y''$ for all $y'' \in Q_i$. Now, there exists $y_{i+1} \in Q_{i+1}$ with $x \succ_r y_{i+1}$ by transitivity and definition of IS. Hence, we can apply above logic to Q_{i+1} as well: $x \succ_r y''$ for all $y'' \in Q_{i+1}$. Inductively, this extends all the way to Q_m , so $x \succ_r y'$ in particular. Since y' is arbitrary, this extends to any $y' \in [y]_r$.

Similar arguments show that $x' \succ_r y$ for any $x' \in [x]_r$. Combining, $x' \succ_r y'$ whenever $x' \in [x]_r$ and $y' \in [y]_r$. \square

Fix a reference point r . Let A_1, \dots, A_n be the distinct equivalence classes of \bowtie_r . By Lemma 5, these sets can be completely ordered by \succ_r , i.e. $A_i \succ_r A_j \iff x \succ_r y$ for all $x \in A_i$ and $y \in A_j$. Label so that $A_1 \succ_r A_2 \succ_r \dots \succ_r A_n$.

Pick an indifference class A_i and an IS Q_1, \dots, Q_M that contains points in every region in A_i . We define $V_i(\cdot)$ on A_i as follows. Define $V_i(x)$ on Q_1 so that $V_i(x) = U^j(x)$ for all $x \in K^j(r)$ where $K^j(r) = Q_1$. Clearly V_i represents \succ_r when restricted to Q_1 . There is no loss in assuming that V_i is bounded, and the closure of its range is an interval.²³

Now, assume inductively that, for a given $m \leq k$, V_i represents \succ_r when restricted to $\bigcup_{j=1}^{m-1} Q_j \equiv \mathbf{Q}^{m-1}$, is bounded, is continuous on \mathbf{Q}^{m-1} , and is an increasing transformation of U^k within Q_j when $Q_j = K^k(r)$. Then, extend V_i to Q_m as follows. By Lemma 5, it is impossible that $y \succ_r x$ for every $x \in \mathbf{Q}^{m-1}$ and every $y \in Q_m$. It will be convenient to relabel regions so that $Q_m = K^m(r)$.

Pick a bounded, strictly increasing, continuous $h : \mathbb{R} \rightarrow \mathbb{R}$. For any $x \in K^m(r)$ so that $x \succ_r y$ for all $y \in \mathbf{Q}^{m-1}$, set

$$V_i(x) = h(U^m(x)) + \beta_+$$

where

$$\beta_+ = \sup\{V_i(x) : x \in \mathbf{Q}^{m-1}\} - \inf\{h(U^m(x)) : x \in K^m(r), x \succ_r y \text{ for all } y \in \mathbf{Q}^{m-1}\}.$$

For any $x \in K^m(r)$ for which there exists $y, y' \in \mathbf{Q}^{m-1}$ so that $y \succ_r x \succ_r y'$, let

$$V_i(x) = \inf\{V_i(y) : y \in \mathbf{Q}^{m-1} \text{ and } y \succsim_r x\}.$$

For all other $x \in K^m(r)$, let

$$V_i(x) = h(U^m(x)) + \beta_-$$

where

$$\beta_- = \inf\{V_i(x) : x \in \mathbf{Q}^{m-1}\} - \sup\{h(U^m(x)) : x \in K^m(r), y \succ_r x \text{ for all } y \in \mathbf{Q}^{m-1}\}.$$

This V_i is bounded and continuous.

²³We can define $V'(x) = h(V(x))$ for $h(v) = -1/(1+v)$ when $v \geq 0$ and $h(v) = -2 + 1/(1-v)$ when $v < 0$.

We now show that it represents \succ_r on \mathbf{Q}^m . Pick $x, y \in \mathbf{Q}^m$. There are four cases:

Case 1: $x, y \in \mathbf{Q}^{m-1}$: then the claim follows by hypothesis.

Case 2: $x \in K^m(r)$ and either $x \succ_r y'$ for all $y' \in \mathbf{Q}^{m-1}$ or $y' \succ_r x$ for all $y' \in \mathbf{Q}^{m-1}$: the claim is immediate.

Case 3: $x \in K^m(r)$ and $y \in \mathbf{Q}^{m-1}$: If $y \succ_r x$, then $y - \epsilon \succ_r x$ for some $\epsilon > 0$ so that $y - \epsilon$ belongs to the same region as y . If $y \sim_r x$, then $V_i(y) \geq V_i(x)$. If this does not hold with equality, then there is a $y' \in \mathbf{Q}^{m-1}$ so that $y' \succ_r x$ and $y \succ_r y'$ (since $y' \not\prec_r y$). But then $y \succ_r x$, a contradiction. If $x \succ_r y$ but $V_i(y) \geq V_i(x)$, there exists $z \in \mathbf{Q}^{m-1}$ so that $V_i(z) \leq V_i(y)$ and $z \succ_r x$. But then by transitivity and hypothesis, $y \succ_r z \succ_r x$.

Case 4: $x, y \in K^m(r)$ and Case 2 does not hold for either x or y : Suppose $x \succ_r y$. If not, then $V_i(y) > V_i(x)$ so there exists a $z \in \mathbf{Q}^{m-1}$ so that $z \succ_r x$ and $z \not\prec_r y$. By weak order, $y \succ_r z$ and so $y \succ_r x$, a contradiction.

Since it represents \succ_r on $K^m(r)$, it also agrees with \succ_m on $K^m(r)$. Hence it is an increasing transformation of U^i within $K^i(r)$ for each $i \leq m$. Renormalize V_i so that its range is a subset of $[-\frac{1}{2} - i, -i]$.

For any $x, y \in A_i$, the above establishes that $V_i(x) \geq V_i(y) \iff x \succ_r y$. For any $x \in A_i$ and $y \in A_j$ where $i < j$, $x \succ_r y$ by Lemma 5 and construction. Since $V_i(x) > -\frac{1}{2} - i$, $V_j(y) < -j$, and $-\frac{1}{2} - i > -j$, we have $V_i(x) > V_j(y)$. Define $U^k(\cdot|r)$ to agree with the appropriate restriction of V_i , and conclude $\{\succ_r\}_{r \in X}$ conforms to CTM under \mathcal{K} . Since r was arbitrary, this completes the proof. \square

A.2. Proof for Theorem 2. Sufficiency is easy to verify. Suppose that $U^k(x) = \sum_{i=1}^n U_i^k(x_i)$. We show that for every category j there exists a vector $w \gg 0$ so that $U^j(x) = \sum_{i=1}^n w_i U_i^k(x_i)$ represents \succ_j on $E^k \cap E^j$.

Consider dimension 1, and the rest follow the same arguments. The goal is to show that $U_1^k(x) - U_1^k(y) \geq U_1^k(a) - U_1^k(b)$ if and only if $U_1^j(x) - U_1^j(y) \geq U_1^j(a) - U_1^j(b)$ for any $x, y, a, b \in E_1^k \cap E_1^j$. If this is the case, then standard uniqueness results give that $U_1^j(x) = \alpha U_1^k(x) + \beta$. The β can be dropped completing the claim.

Let π_i be the projection onto the i -coordinate. Then, $E_1^k = \pi_1(E^k)$ is open and connected for any category k . This follows from E^k connected and open and π_i continuous. In \mathbb{R} , connected implies convex.

Claim 1. *For any $z \in E_1^k \cap E_1^j$, there exists a neighborhood $O_z = B_{\epsilon_z}(z)$ so that $U_1^k(x) - U_1^k(y) \geq U_1^k(a) - U_1^k(b)$ if and only if $U_1^j(x) - U_1^j(y) \geq U_1^j(a) - U_1^j(b)$ for any $x, y, a, b \in O_z$.*

To see it is true, pick $x \in E_1^k \cap E_1^j$. Then there is an $a^l \in E^l$ with $a_1^l = x$ for $l = k, j$. Let $U_{-i}^k(y) = \sum_{j \neq i} U_j^k(y_j)$ for any $y \in X$. Since each $a^l \in K^l(r^l)$ for some

$r^l \in X$, there exists an $\epsilon^l > 0$ so that $B_{2\epsilon^l}(a^l) \subset K^l(r^l) \subset E^l$, where the distance is given by the supnorm. Pick $\epsilon \in (0, \epsilon^l)$ so that

$$U_1^l(x + \epsilon) - U_1^l(x - \epsilon) < U_{-1}^l(a^l + \epsilon^l) - U_{-1}^l(a^l - \epsilon^l)$$

for $l = k, j$. Then, for any $a, b \in [x - \epsilon, x + \epsilon]$ there exists y_{-1}^a, y_{-1}^b so that $(a, y_{-1}^a), (b, y_{-1}^b) \in B_{2\epsilon^k}(a^k)$ and $(a, y_{-1}^a) \sim_{r^k} (b, y_{-1}^b)$ by Category Continuity and CM. In particular, $U_1^k(a) - U_1^k(b) = U_{-1}^k(y_{-1}^b) - U_{-1}^k(y_{-1}^a)$. For any $a', b' \in [x - \epsilon, x + \epsilon]$, it holds that $U_1^k(a) - U_1^k(b) \geq U_1^k(a') - U_1^k(b')$ if and only if $(b', y_{-1}^a) \succsim_{r^k} (a', y_{-1}^b)$. Similarly, there exist z_{-1}^a, z_{-1}^b so that $(a, z_{-1}^a), (b, z_{-1}^b) \in B_{2\epsilon^j}(a^j)$ and $(a, z_{-1}^a) \sim_{r^j} (b, z_{-1}^b)$. Now, $(b', z_{-1}^b) \succsim_{r^j} (a', z_{-1}^a)$ if and only if $U_1^j(a) - U_1^j(b) \geq U_1^j(a') - U_1^j(b')$. By Reference Interlocking and weak order, $(b', z_{-1}^b) \succsim_{r^j} (a', z_{-1}^a)$ if and only if $(b', y_{-1}^a) \succsim_{r^k} (a', y_{-1}^b)$, so we conclude that the claim holds with $\epsilon_x = \epsilon$.

We now extend to the entire domain (this follows similar arguments in CW). Pick an arbitrary $x_* < x^* \in E_1^k \cap E_1^j$ and consider $Z = (x_*, x^*]$. If the claim is true, then standard uniqueness results give that $U_1^j(x) = \alpha U_1^k(x) + \beta$ for all $x \in O_z$ for some $\alpha > 0$. Let α^*, β^* be the constants so that $U_1^j(x) = \alpha^* U_1^k(x) + \beta^*$ for all x in the neighborhood of x^* , as guaranteed to exist by the claim.

Let

$$Z_1 = \{s \in Z : U_1^j(x) = \alpha^* U_1^k(x) + \beta^* \text{ for all } x \in (x_*, s]\}.$$

Z_1 is not empty by the claim. We show that it is both open and closed by picking any $s_1 \in cl(Z_1)$ and showing $s_1 \in int(Z_1)$. Since $[x_*, s_1]$ is compact and $O = \{O_z : z \in [x_*, s_1]\}$ is an open covering, there exists $\{O_1, \dots, O_n\} \subset O$ with $x_* \in O_1$, $s_1 \in O_n$ and $O_m \cap O_{m'} = \emptyset$ for all $m' \geq m + 2$. On each O_m , there exists α_m, β_m so that the utility indexes agree by the claim. Also, O_m and O_{m+1} have non-empty intersections with more than two points, so $(\alpha_{m+1}, \beta_{m+1}) = (\alpha_m, \beta_m)$. In particular, O_1 intersects O_{x_*} so $\alpha_m = \alpha^*$ for all m . Then $O_n \cap Z \subset Z_1$, i.e. $s_1 \in int(Z_1)$, so $cl(Z_1) \subset int(Z_1) \subset Z_1 \subset cl(Z_1)$, i.e. Z_1 is both closed and open relative to Z . Conclude $Z_1 = Z$ since Z connected.

Since $U_1^j(x) = \alpha U_1^k(x) + \beta$ for all $x \in (x_*, x^*]$ for any interval in the domain, it holds for the whole domain as well. Extend to other categories that intersect $E_1^i \cup E_1^j$ inductively. If there is no intersecting category, we can start again and obtain a (dis-joint) interval, the values of U_1^i (and U_1^j) on which have no bearing on the DM's choices. Similar arguments obtain for the other dimensions. Moreover, there is no loss in setting each $\beta = 0$. This completes the proof. \square

A.3. Proof of Theorem 3. To save notation, until after Lemma 10, we fix r and write K^k instead of $K^k(r)$ and \succsim instead of \succsim_r . We also identify $x\alpha^k y$ with the alternative $\alpha x \oplus^k (1 - \alpha)y$. Let (U^1, \dots, U^n) be the additive functions that represent

$\succsim_1, \dots, \succsim_n$. Observe that $U^k(x\alpha^k y) = \alpha U^k(x) + (1 - \alpha)U^k(y)$ for any α , provided that $x, y, x\alpha^k y \in E^k$.

Recall from Definition 6 that an indifference sequence is a finite sequence of categories with indifference between each succeeding members.

Definition 7. The function v is a *utility for the indifference sequence* (Q_1, \dots, Q_m) if v is an increasing additive utility function on each Q_k and for all $k, x, y \in Q_k \cup Q_{k+1}$: $x \succsim y \iff v(x) \geq v(y)$.

Lemma 6. If $x^k \in K^k$, $x^l \in K^l$, and $x^k \sim x^l$, then there is $a > 0, b \in \mathbb{R}$ such that for $x \in K^k$ and $y \in K^l$, $x \succsim y \iff U^k(x) \geq \alpha U^l(y) + \beta$.

Proof. W.L.O.G., take $U^k(x^k) = 0$. There is $\epsilon_k > 0$ such that $B_{2\epsilon_k}(x^k) \subset K^k$. By CM and Category Continuity, there is $\epsilon_l > 0$ such that $B_{\epsilon_l}(x^l) \subset K^l$ and for all $y \in B_{\epsilon_l}(x^l)$, $x^* = x^k + \epsilon_k \succ y \succ x^k - \epsilon_k = x_*$. For any $y \in K^l$ and α such that $y\alpha^l x^l \in B_{\epsilon_l}(x^l)$, there exists $\beta \in (0, 1)$ such that $x^* \beta^k x_* \sim y\alpha^l x^l$ by Category Continuity, CM, and that \succsim is a weak order. Let $V^l(y) = \alpha^{-1}U^k(x^* \beta^k x_*)$. This is well defined, additive, increasing, and ranks alternatives in the same way as U^l . Thus, $V^l(y) = aU^l(y) + b$ for some $a > 0$ and $b \in \mathbb{R}$.

For any $x \in K^k$ and $y \in K^l$, pick $\alpha \in [0, 1]$ such that $x\alpha^k x^k \in B_{\epsilon_k}(x^k)$ and $y\alpha^l x^l \in B_{\epsilon_l}(x^l)$. By construction, $y\alpha^l x^l \sim y'$ when $y' \in B_{\epsilon_k}(x^k)$ and $U^k(y') = \alpha V_l(y)$. Thus, $x\alpha^k x^k \succsim y' \sim y\alpha^l x^l$ holds if and only if $U^k(x) \geq V_l(y)$ and $x \succsim y \iff x\alpha^k x^k \succsim y\alpha^l x^l$ by AAC since $x^k \sim x^l$, completing the proof. \square

For an indifference sequence (Q_1, \dots, Q_m) with utility v , we label the range of utilities as $cl(v(Q_k)) = [l_k, u_k]$ where $l_k \leq u_k$. Note that we allow $Q_k = Q_l$ for $k \neq l$.

Lemma 7. For an indifference sequence (Q_1, \dots, Q_m) , there is an affine, increasing utility v for it.

Proof. The proof is by induction. We claim that there is a utility $v^k : X \rightarrow \mathbb{R}$ that is a utility for the IS (Q_1, \dots, Q_k) for any k . When $k = 1$ or $k = 2$, this is true by the above lemmas. The induction hypothesis (IH) is that the claim is true for $k = N$. Consider $k = N + 1$. Let v^N be the utility for (Q_1, \dots, Q_N) be index that exists by the IH. If $Q_{N+1} \subseteq \bigcup_{i=1}^N Q_i$, then we are done. If not, then for $Q_N = K^l$, there is no loss in normalizing v^N so that it equals U^l on $K^l(r)$. Suppose $Q_{N+1} = K^j(r)$, and let α, β be the scalars claimed to exist by Lemma 6, so that $U^j(x) \geq \alpha U^l(y) + \beta \iff x \succsim_r y$ for $x \in K^k(r)$ and $y \in K^l(r)$. Restricted to Q_N , $v^N = U^l$, so we can define $v^{N+1}(x) = \alpha v^N(x) + \beta$ if $x \in \bigcup_{i=1}^N Q_i$ and

$$v^{N+1}(x) = U^j(x)$$

if $x \in Q_{N+1}$. Then, if $l < N$ and $x, y \in Q_l \cup Q_{l+1}$, then we are done by the IH, since $v^{N+1}(x) \geq v^{N+1}(y) \iff v^N(x) \geq v^N(y)$. If $x, y \in Q_N \cup Q_{N+1}$, then Lemma 6 and construction implies the result. The claim then holds by induction. \square

Lemma 8. *Fix an indifference sequence (Q_1, \dots, Q_n) with utility v . If $x^k \in Q_k$ for $k = i, i+1, i+2$ with $x^i \sim x^{i+1} \sim x^{i+2}$, then $(Q_1, \dots, Q_i, Q_{i+2}, \dots, Q_n)$ is an indifference sequence (after relabeling) with utility v .*

Proof. The Lemma is vacuously true for any 1 or 2-element IS. Fix an IS (Q_1, \dots, Q_n) with $n \geq 3$ and v as above, and suppose $x^k \in Q_k$ for $k = i, i+1, i+2$ with $x^i \sim x^{i+1} \sim x^{i+2}$. By transitivity $x^i \sim x^{i+2}$, so $(Q_1, \dots, Q_i, Q_{i+2}, \dots, Q_n)$ is an IS; it remains to be shown that v is a utility for it. There is an $\epsilon > 0$ s.t. $B = B_\epsilon(v(x^i)) \subset (l_k, u_k)$ for $k = i, i+1, i+2$. Let $v^{-1}(u) : B \rightarrow Q_{i+1}$ be an arbitrary point in Q_{i+1} such that $v[v^{-1}(u)] = u$. Now, fix $x \in Q_i$ and $y \in Q_{i+2}$. For α small enough, $v(x\alpha^i x^i), v(y\alpha^{i+2} x^{i+2}) \in B$. Then $x\alpha^i x^i \sim v^{-1}(v(x\alpha^i x^i))$ and $y\alpha^{i+2} x^{i+2} \sim v^{-1}(v(y\alpha^{i+2} x^{i+2}))$. So

$$\begin{aligned} x \succsim y &\iff x\alpha^i x^i \succsim y\alpha^{i+2} x^{i+2} \\ &\iff v^{-1}(v(x\alpha^i x^i)) \succsim v^{-1}(v(y\alpha^{i+2} x^{i+2})) \\ &\iff v[v^{-1}(v(x\alpha^i x^i))] \geq v[v^{-1}(v(y\alpha^{i+2} x^{i+2}))] \\ &\iff \alpha v(x) + (1 - \alpha)v(x^i) \geq \alpha v(y) + (1 - \alpha)v(x^{i+2}) \\ &\iff v(x) \geq v(y) \end{aligned}$$

This establishes the Lemma. \square

Lemma 9. *Fix an indifference sequence (Q_1, \dots, Q_n) with utility v . If $(l_1, u_1) \cap (l_n, u_n) \neq \emptyset$, then there exists i and $x^k \in Q_k$ for $k = i, i+1, i+2$ with $x^i \sim x^{i+1} \sim x^{i+2}$.*

Proof. If there is i with $(l_i, u_i) \cap (l_{i+2}, u_{i+2}) \neq \emptyset$, then there is $u \in \bigcap_{j=i, i+1, i+2} (l_j, u_j)$ so there exists $x_j \in Q_j$ with $v(x_j) = u$ for $j = i, i+1, i+2$ and thus by the hypothesis, $x_i \sim x_{i+1} \sim x_{i+2}$. We show there exists such an i by contradiction. If $l_{i+2} > u_i$ for all i or $l_i > u_{i+2}$ for all i , then $(l_1, u_1) \cap (l_n, u_n) = \emptyset$, a contradiction. So there must exist i such that $[l_{i+2} > u_i \text{ and } l_{i+2} > u_{i+4}]$ or $[u_{i+2} < l_i \text{ and } u_{i+2} < l_{i+4}]$. In the first case, $l_{i+2} \in (l_{i+1}, u_{i+1}) \cap (l_{i+3}, u_{i+3})$; in the second, $u_{i+2} \in (l_{i+1}, u_{i+1}) \cap (l_{i+3}, u_{i+3})$. In either case, we have a contradiction. \square

Lemma 10. *Fix an indifference sequence (Q_1, \dots, Q_n) with utility v . Then for all $x, y \in \bigcup_i Q_i$, $x \succsim y \iff v(x) \geq v(y)$.*

Proof. This is clearly true if $n = 1$. (IH) Suppose the claim is true for any IS with $m < n$ elements. Fix an IS (Q_1, \dots, Q_n) with utility v . If $x \notin Q_1 \cup Q_n$ or $y \notin Q_1 \cup Q_n$, then the claim immediately follows from the IH, and clearly holds if $x, y \in Q_i$ for some i . So it suffices to consider arbitrary $x \in Q_1$ and $y \in Q_n$. By Lemmas 8 and 9, if $(u_1, l_1) \cap (l_n, u_n) \neq \emptyset$, we can form a shorter IS from Q_1 to Q_n and the claim then follows from the IH.

There are two cases to consider: $l_n > u_1$ and $u_n < l_1$. Consider $l_n > u_1$. The range of v restricted to $\bigcup_{i=1}^{n-1} Q_i$ is dense in $\bigcup_{i=1}^{n-1} (l_i, u_i) = (\bar{l}, \bar{u})$. Note $l_n \in (\bar{l}, \bar{u})$ since $x_{n-1} \sim y_n$, so $(l_{n-1}, u_{n-1}) \cap (l_n, u_n) \neq \emptyset$. Then $(l_n, v(y))$ is an open interval having a non-empty intersection with (\bar{l}, \bar{u}) . Since the range of v is dense in (\bar{l}, \bar{u}) , there exists $y' \in Q_{n'}$ with $l_n < v(y') < v(y)$. Since $l_n > u_1$, $n' > 1$. Then $(Q_1, \dots, Q_{n'})$ and $(Q_{n'}, \dots, Q_n)$ are both ISes with strictly less than n elements. Applying the IH, $y' \succ x$ and $y \succ y'$. Conclude using transitivity that $y \succ x$. Similar arguments obtain the desired conclusion when $u_n < l_1$. \square

Define \bowtie_r as in the proof of Theorem 1, and let A_1, \dots, A_n be the distinct indifference classes of \bowtie_r . Again using Lemma 5, we can relabel so that $x \in A_i$ and $y \in A_{i+1}$ implies $x \succ_r y$. By Lemma 10, there is v_i on A_i so that v_i is additive and increasing within categories and $x \lesssim_r y \iff v_i(x) \geq v_i(y)$ for all $x, y \in A_i$.

By Unbounded and Lemma 5, every positive unbounded region (if any) is a subset of A_1 , and every negative unbounded region (if any) is a subset of A_n . If one region is both positive and negative unbounded, then $n = 1$. Therefore, $v_i(A_i)$ is bounded for all $i \in (1, n)$, and $v_n(A_n)$ is bounded above whenever $n > 1$. Define $V(x) = v_1(x)$ for all $x \in A_1$. For $x \in A_i$ with $i > 1$, define $V(x)$ recursively by

$$V(x) = v_i(x) - \sup_{y \in A_i} v_i(y) + \inf_{y \in A_{i-1}} V(y) - 1.$$

Observe $V(\cdot)$ is a positive affine transformation of $v_i(\cdot)$ when restricted to A_i , and if $x \in A_i$, $y \in A_j$ and $i > j$, then $V(x) > V(y)$. Thus V represents \lesssim_r and, when restricted to any given region, is affine and increasing.

Defining $U^k(\cdot|r)$ as the (unique) affine transformation of U^k so it agrees with V on $K^k(r)$ establishes that \lesssim_r is an affine CTM. Since r was arbitrary, this establishes that each \lesssim_r has such a representation. Conclude that $\{\lesssim_r\}$ conforms to Affine CTM, completing the proof. \square

A.4. Proof of Theorem 4. Without loss of generality, normalize so that $U^1(\cdot|r) = U^1(\cdot|r')$ for all r, r' . Suppose $U^k(\cdot|r) \neq U^k(\cdot|r')$ for some r, r' and some k . Then, let $\bar{\epsilon} = d(r, r')$ and pick a sequence $\hat{r}_n \rightarrow \hat{r}$ such that: $U^k(\cdot|\hat{r}_n) \neq U^k(\cdot|r)$, $\hat{r}_n \in B_{\bar{\epsilon}}(r)$ for all n , and $d(\hat{r}_n, r) \rightarrow \inf\{d(r', r) : U^k(\cdot|r) \neq U^k(\cdot|r')\}$. Since $\hat{r}_n \in cl(B_{\bar{\epsilon}}(r))$, there is no loss in assuming this sequence converges. Similarly, let r_n be a sequence in $B_{\bar{\epsilon}}(r)$ such that $r_n \rightarrow \hat{r}$ and $U^k(\cdot|r) = U^k(\cdot|r_n)$.

By hypothesis and that each $K^k(r)$ is open, there exists $\epsilon > 0$, x^k and x^1 such that $B_{2\epsilon}(x^k) \subset K^k(\hat{r})$, $B_{2\epsilon}(x^1) \subset K^1(\hat{r})$, and $x^k \sim_{\hat{r}} x^1$. By continuity of the region functions, $B_{\epsilon}(x^k) \subseteq K^i(\hat{r}_n) \cap K^i(r_n)$ and $B_{\epsilon}(x^1) \subseteq K^1(\hat{r}_n) \cap K^1(r_n)$ for n large enough. For z close enough to x^k , there exists $y(z) \in B_{\epsilon}(x^1)$ such that $z \sim_{\hat{r}} y(z)$. But then by SC, $z \sim_{r_n} y(z)$ and $z \sim_{\hat{r}_n} y(z)$. Thus $U^k(z|r_n) = U^1(y(z)|r_n) = U^1(y(z)|\hat{r}_n) = U^k(z|\hat{r}_n)$ for

all z close enough to x_k , implying that $U^k(\cdot|r_n) = U^k(\cdot|\hat{r}_n)$, a contradiction. Conclude $U^k(\cdot|r) = U^k(\cdot|r')$ for all r, r' . \square

A.5. Examples from Table 1. Example 1 shows that BGS violates Cancellation and inspecting Figure 1 shows it violates Monotonicity. It remains to show that TK violates Reference Irrelevance and that MO violates Cancellation. This is established by the following two examples.

Example 2 (TK violates Reference Irrelevance). Consider a TK model with $\lambda_1 = \lambda_2 = 2$. Then, for $r = (10, 10)$, $x = (12, 12)$ and $y = (9, 16)$, $y \succsim_r x$ since $(12 - 10) + (12 - 10) = 2(9 - 10) + (16 - 10)$. For $r' = (11, 11)$, $x \succ_r y$ since $(12 - 11) + (12 - 11) > 2(9 - 11) + (16 - 11)$. But $x \in R_1^{GL}(r) \cap R_1^{GL}(r')$ and $r \in R_2^{GL}(r) \cap R_2^{GL}(r')$, so the family violates Reference Irrelevance.

Example 3 (MO violates Cancellation). Let $Q(r) = \{x \in X : x_1/2 + x_2 > r_1/2 + r_2\}$ and $c(r) = 1$. Then, let $x = (2, 1)$, $y = (1, 2)$, $z = (4, 4)$, and $r = (0.9, 1.9)$. Since $(x_1, z_2) = (2, 4) \succsim_r (4, 2) = (z_1, y_2)$ and $(z_1, x_2) = (4, 1) \succsim_r (1, 4) = (y_1, z_2)$ because all four points belong to $Q(r)$, cancellation requires that $x \succsim_r y$. However, $x \notin Q(r)$, so $y \succ_r x$, so cancellation does not hold.

A.6. Other models and CTM. In this subsection, we present the functional forms of the other models of salience we discussed, and show that they are not CTM.

- Gabaix [2014] assumes a rational DM would maximize $u(a, w)$ but actually maximizes

$$u(a, (w_1 m_1^*, \dots, w_n m_n^*))$$

where

$$m^* \in \arg \min_{m \in [0,1]^n} \frac{1}{2} \sum_{i,j} (1 - m_i) \Lambda_{ij} (1 - m_j) + \kappa \sum_i m_i^\alpha$$

where Λ_{ij} incorporates the “variance” in the marginal utility of dimensions i and j . When n is large, m_i^* is often zero, so $(w_1 m_1^*, \dots, w_n m_n^*)$ is a “sparse” vector.

- Tversky & Kahneman [1991] refer in general to

$$V_{CTK}(x|r) = \sum_i v_i(u_i(x_i) - u_i(r_i))$$

where v_i is concave above 0 and convex below

- Bordalo et al. [2019] and the continuous form of the salient thinking model has

$$V_{CBGS}(x|r) = w(x_1, r_1)x_1 + w(x_2, r_2)x_2$$

where w has the same properties as a salience function.

- Munro & Sugden [2003] use the functional from

$$V_{MS}(x|r) = A(r) \left(\sum_i \gamma_i r_i^{\rho-\beta} x_i^\beta \right)^{\frac{1}{\beta}}$$

- Bhatia & Golman [2013] assume that the DM chooses the bundle x that maximizes

$$U(x|r) = \alpha_1(r_1)[V(x_1) - V(r_1)] + \alpha_2(r_2)[V(x_2) - V(r_2)]$$

given that a reference point r , where each α_i is increasing and positive.

The first fails to be CTM, as the indifference curves have the same slope everywhere for a fixed context. If they were CTM, then they would necessarily have only a single region. Single region CTM coincides with the neoclassical model. The final four explicitly take into account a reference point. In all four, it is easy to see that the reference point affects the marginal rate of substitution between attributes. This implies a violation of weak reference irrelevance for any given category function: any two points in the same category that are indifferent to each other necessarily remain so for a sufficiently small change in the reference point.

A.7. Proof of Proposition 2. K satisfying S0-S4 implies that $E^1 = E^2 = \mathbb{R}_{++}^n$, so the structure assumption is satisfied. Moreover, Theorem 5 gives that the categories are generated by a salience function. The axioms allow us to apply Theorems 2 and 4 to get a strong CTM representation of the family with reweighted utility indexes. Hence,

$$U^k(x) = w_1^k u_1(x_1) + w_2^k u_2(x_2) + \beta^k$$

for each $x \in X$.

There is no loss in normalizing so that $\beta^1 = 0$. Pick $x \in X$ with $x_1 > x_2$, and by S4 observe that $x \in K^1(r)$ for $r = (x_1, x_2/2)$ and $x \in K^2(r')$ for $r' = (x_1/2, x_2)$. Since $K^1(r)$ and $K^2(r')$ are open, there exists $\epsilon > 0$ so that $B_\epsilon(x) \subset K^1(r) \cap K^2(r')$. Since U^1 is continuous and increasing, there is $y \in B_\epsilon(x)$ with $y_1 < x_1$ so that $U^1(y) = U^1(x)$, i.e. $y \sim_r x$; this y necessarily has $y_2 > x_2$ by CM. Then, SDO implies $y \succ_{r'} x$, i.e. $U^2(y) > U^2(x)$, which requires $w_2^1/w_2^2 < w_1^1 > w_1^2$. We can incorporate β^2 into u_2 by replacing it with $u_2 + \beta^2/(w_2^2 - w_2^1)$ or into u_1 by replacing β^2 into u_1 by replacing it with $u_1 + \beta^2/(w_1^2 - w_1^1)$. At least one does not involve dividing by zero, as otherwise $w_i^2 = w_i^1$ for $i = 1, 2$. \square

A.8. TK. This subsection states and proves a characterization theorem for TK.

Proposition 5. *A family of preferences $\{\succsim_r\}_{r \in X}$ has a TK representation if and only if it is an affine CTM with a gain-loss regional function that satisfies Reference Interlocking, Monotonicity, Cancellation, and continuity of each \succsim_r .*

Tversky & Kahneman [1991, p. 1053] provide an alternative axiomatic characterization of the model, and our result makes heavy use of their theorem.

Proof. Necessity follows from the discussion above and TK's theorem. To show sufficiency, we rely on TK's theorem, which states that any monotone, continuous family of preference relations that satisfies cancellation, sign-dependence and reference interlocking has a TK representation. Given our assumptions, we need to show that $\{\succsim_r\}$ satisfies sign-dependence and reference interlocking.

TK say that $\{\succsim_r\}$ satisfies sign-dependence if “for any $x, y, r, s \in X$, $x \succsim_r y \iff x \succsim_s y$ whenever x and y belong to the same quadrant with respect to r and with respect to s , and r and s belong to the same quadrant with respect to x and with respect to y .” This happens if and only if $x \in K^k(r) \cap K^k(s)$ and $y \in K^k(r) \cap K^k(s)$ for some $k \in \{1, 2, 3, 4\}$. Then, sign-dependence is exactly an implication of affine CTM, since $U^k(\cdot|r) = \alpha U^k(\cdot|s) + \beta$ for $\alpha > 0$.

TK say that $\{\succsim_r\}$ satisfies reference interlocking if “for any $w, w', x, x', y, y', z, z'$ that belong to the same quadrant with respect to r as well as with respect to s , $w_1 = w'_1, x_1 = x'_1, y_1 = y'_1, z_1 = z'_1$ and $x_2 = z_2, w_2 = y_2, x'_2 = z'_2, w'_2 = y'_2$, if $w \sim_r x$, $y \sim_r z$, and $w' \sim_s x'$ then $y' \sim_s z'$.” The assumptions on quadrants imply that $w, w', x, x', y, y', z, z' \in K^k(r) \cap K^l(s)$ for some $k, l \in \{1, 2, 3, 4\}$. Since $y', z' \in K^l(s)$, the conclusion follows immediately from RI. \square

A.9. Example 4.

Example 4. The following salience functions generates regions all satisfy S0-S3, but only satisfy a subset of the other properties.

- (1) The function $\sigma(x, r) = \frac{\max\{x, r\}^2}{\min\{x, r\}}$ generates regions that violate S4-S6. Note $\sigma(a, a) = a$ for $a > 0$. Then $(a, b + \epsilon), (a, b) \in K^1(a, b)$ for all $a > b$ and small enough $\epsilon > 0$, contradicting S4 and S6, respectively. Also note $\sigma(a, a) = \sigma(\sqrt{a}, 1)$ for $a > 0$. Hence, $(a, \sqrt{a}) \notin K^1(a, 1)$ but $(a + \epsilon, \sqrt{a}) \in K^1(a + \epsilon, 1)$ for every $\epsilon > 0$, violating S5.
- (2) The function $\sigma(x, r) = |x^2 - r^2|$ generates regions that satisfy S0-S4 but violate S5 and S6. Observe that $(2, \sqrt{5}) \notin K^1(1, \sqrt{2})$ since $\sigma(2, 1) = \sigma(\sqrt{5}, \sqrt{2}) = 3$, but $(2 + \epsilon, \sqrt{5}) \in K^1(1 + \epsilon, \sqrt{2})$ for any $\epsilon > 0$ since $\sigma(2 + \epsilon, 1 + \epsilon) = 3 + 2\epsilon > 3$, contradicting S5. It is routine to verify S4 by differentiating. Also, $x = (2, 2)$ and $r = (4, 1)$ have $x_1 x_2 = r_1 r_2$, but $\sigma(2, 4) > \sigma(2, 1)$, so $x \in K^1(r)$, contradicting S6.
- (3) The function $\sigma(x, r) = |\sqrt{x} - \sqrt{r}|$ generates regions that satisfy S0-S5 but violate S6. Also, $x = (2, 2)$ and $r = (4, 1)$ have $x_1 x_2 = r_1 r_2$, but $\sigma(2, 4) > \sigma(2, 1)$, so $x \in K^1(r)$, contradicting S6. Differentiating establishes S4 and S5.

- (4) The function $\sigma(x, r) = \frac{\max\{x, r\}}{\min\{x, r\}}$ generates regions that satisfy S0-S6.

A.10. Proof of Theorem 5. We first prove the following lemma.

Lemma 11. *If \mathcal{K} is a category function, then for any $\epsilon > 0$ and x so that $B_\epsilon(x) \subset K^i(r)$, there exists $\delta > 0$ so that $B_{\epsilon/2}(x) \subset K^i(r')$ for all $r' \in B_\delta(r)$.*

Proof. Let $B_{\epsilon/2}(x)$. For each $j \neq i$, $d(K^j(r), B) > \epsilon/2$, where $d(\cdot)$ is the Hausdorff metric,²⁴ and continuity of K^j implies that there exists a neighborhood O_j of r so that $d(K^j(r'), B) > \epsilon/4$ for all $r' \in O_j$. Let $O = \bigcap_{j \neq i} O_j$. Then, for any $r' \in O$, $B \notin cl(\bigcup_{j \neq i} K^j(r'))$. Since $cl(\bigcup_i K^i(r')) = X$, $B \subset cl(K^i(r'))$. But since B is open, $B \subset int(cl(K^i(r')))$ since $K^i(r')$ is regular open. \square

For sufficiency, define a binary relation S by $(a, b)S(c, d)$ if and only if $(a, c) \notin K^2(b, d)$. S is clearly complete. It is also transitive by S3. We show it has an open contour sets. Let S^* be the strict part of S . If $(a, b)S^*(c, d)$, then $x \in K^1(r)$ for $x = (a, c)$ and $r = (b, d)$. $K^1(r)$ is open by S0 so there exists $\epsilon > 0$ so that $B_\epsilon(x) \subset K^1(r)$. By Lemma 11, $x \in K^1(r')$ for all r' in a neighborhood O' of r . Conclude $(a', b')S^*(c', d')$ for all $(a', b'), (c', d') \in B_\epsilon(x) \times O'$. Standard results then show existence of a continuous function σ so that $(a, b)S(c, d)$ if and only if $\sigma(a, b) \geq \sigma(c, d)$. σ is symmetric by S2 and increasing in contrast by S1 and S4. Hence $x \in K^1(y)$ if and only if $\sigma(x_1, y_1) > \sigma(x_2, y_2)$, and by S2, $x \in K^2(y)$ if and only if $y' \in K^1(x')$ where x', y' are the reflections of x, y . Hence, $x \in K^2(y)$ if and only if $\sigma(x_1, y_1) < \sigma(x_2, y_2)$.

Pick any a, b . By S3, $\sigma(a, b) = \sigma(b, a)$ so $(a, b) \notin K^1(b, a)$ for any a, b . By S5, $(a + \epsilon, b) \notin K^1(b + \epsilon, a)$. Then, $(b, a)S(a + \epsilon, b + \epsilon)$ so $\sigma(a, b) = \sigma(b, a) \geq \sigma(a + \epsilon, b + \epsilon)$. Since a, b were arbitrary, diminishing sensitivity holds.

For necessity, verifying that S0-S5 hold are trivial, except that each $K^i(r)$ is regular open. To see this, pick r and $x \in int(cl(K^1(r)))$ (symmetric arguments hold for K^2). Suppose $x \gg r$ (the other cases follow by changing the signs). Then, there is $\epsilon > 0$ so that $\bar{x} = (x_1 - \epsilon, x_2 + \epsilon) \in cl(K^1(r))$. Then, there exists $x' \in K^1(r)$ that is arbitrarily close to \bar{x} , and we can take it so that $r_1 < x'_1 < x_1$ and $r_2 < x'_2 < x_2$. Then, $\sigma(x_1, r_1) > \sigma(x'_1, r_1) > \sigma(x'_2, r_2) > \sigma(x_2, r_2)$ since σ is increasing in contrast. Hence $x \in K^1(r)$ and since x was arbitrary, $int(cl(K^1(r))) \subset K^1(r)$. Clearly, $K^1(r) \subset int(cl(K^1(r)))$.

Now we show the following are equivalent:

- (i) The functions K^1 and K^2 satisfy S0, S1, and S6,
- (ii) There exists a salience function σ s.t. $x \in K^k(r) \iff \sigma(x_k, r_k) > \sigma(x_{-k}, r_{-k})$

²⁴In this case it is actually a pseudo metric.

That (ii) implies (i) follows from the first part, and that S6 is implied by symmetry and homogeneity of degree zero. Now, we show (i) implies (ii). Set $\sigma(a, b) = \max\{a/b, b/a\}$. Clearly σ is a salience function, and we show that σ generates K^1 and K^2 . Fix $r \in X$ and set $A = \{x : \sigma(x_1, r_1) > \sigma(x_2, r_2)\}$. We show $A = K^1(r)$.

Claim $A \cap K^2(r) = \emptyset$. If not, pick $x \in A \cap K^2(r)$. $x \in A$ implies either (a) $x_1/r_1 > x_2/r_2$ and $x_1/r_1 > r_2/x_2$ or (b) $r_1/x_1 > x_2/r_2$ and $r_1/x_1 > r_2/x_2$. If (a) and $x_2 \leq r_2$, then

$$x_1/r_1 > r_2/x_2 \geq x_2/r_2 \text{ implies } x_1 > r_1 r_2 / x_2 \geq r_1,$$

so there exists $\lambda \in [0, 1)$ such that $(\lambda x_1 + (1 - \lambda)r_1, x_2) = (r_1 r_2 / x_2, x_2) = x'$. If (a) and $x_2 > r_2$, then

$$x_1 > r_1 x_2 / r_2 > r_1,$$

so there exists $\lambda \in (0, 1)$ such that $(\lambda x_1 + (1 - \lambda)r_1, x_2) = (r_1 x_2 / r_2, x_2) = x'$. By S1 and $x \in K^2(r)$, $x' \in K^2(r)$. However, we have either $x'_1 x'_2 = r_1 r_2$ or $x'_1 / x'_2 = r_1 / r_2$ so $x' \notin K^2(r)$ by S6, a contradiction. A similar contradiction obtains if (b) holds.

Now, since $A \cap K^2(r) = \emptyset$ and $K^1(r) \cup K^2(r)$ is dense, $A \subset cl(K^1(r))$. By S0, $K^1(r) = int(cl(K^1(r)))$. Since A is an open set contained in $cl(K^1(r))$, $A \subseteq K^1(r)$. Similarly, for $B = \{x : \sigma(x_1, r_1) < \sigma(x_2, r_2)\}$, $B \subseteq K^2(r)$. But

$$(A \cup B)^c = \{x : x_1 x_2 = r_1 r_2 \text{ or } x_1 / x_2 = r_1 / r_2\},$$

and by S6, $(A \cup B)^c \cap K^k(r) = \emptyset$ for $k = 1, 2$. Thus $A = K^1(r)$ and $B = K^2(r)$, completing the proof.

Finally, fix any HOD salience function s . Observe $s(a, b) > s(c, d)$ if and only if $s(a/b, 1) > s(c/d, 1)$ by homogeneity if and only if $s(\max(a/b, b/a), 1) > s(\max(c/d, d/c), 1)$ by symmetry if and only if $\max(a/b, b/a) > \max(c/d, d/c)$ by ordering. Thus if one salience function generates the regions, every other salience function does as well. \square

A.11. Proof of Proposition 1. Pick any $r \in X$. Observe that $x = (r_1 + k, r_2)$ necessarily belongs to $K^1(r)$ by S4, as does an open set $O \ni x$. This set O can be identified by looking at whether Cancellation and Monotonicity hold on the set. By varying r and k , we obtain a covering of the entirety of \mathbb{R}_{++}^2 by points that necessarily belong to the 1-salient region. This allows one to identify \succsim^1 and obtain a representation U^1 . Repeating with $x' = (r_1, r'_2 + k)$ obtains a representation U^2 of \succsim^2 .

Fix any $r \in X$. Consider $y \gg r$ and let

$$I^*(y) = \{y' : U^k(y') = U^k(y) \text{ and } U^{-k}(y) \neq U^{-k}(y')\} \setminus \{y\}.$$

If $y \in K^k(r)$, then there exists $y' \in I^*(y)$ arbitrarily close to y so that $y, y' \in K^k(r)$; for any such y' , $y \sim_r y'$. If $y' \not\sim_r y$ for every $y' \in I^*(y) \cap B_\epsilon(y)$ for some $\epsilon > 0$, $y \notin K^k(r)$.

Conclude $y \in K^k(r)$ if and only if there exists $y' \in I^*(y) \setminus \{y\}$ arbitrarily close to y so that $y \sim_r y'$.

We now infer whether $\sigma(x, a) > \sigma(y, b)$ for any x, y, a, b by considering whether an alternative x' is in $K^1(r)$ for appropriately chosen bundles so that $x' \gg r$. This is impossible if $x = a$ and always true if $y = b$ and $x \neq a$. For any other values, we have that $\sigma(x, a) > \sigma(y, b)$ if and only if either $(x, y) \in K^1(a, b)$, $x > a$ and $y > b$; $(x, b) \in K^1(a, y)$, $x > a$, and $b > y$; $(a, y) \in K^1(x, b)$, $x < a$, and $y > b$; or $(a, b) \in K^1(x, y)$, $x < a$, and $b > y$. In this way we can reveal the σ function and thus the regions. \square

APPENDIX B. PROOFS AND EXTRAS FROM SECTION 5

B.1. Axioms for c . This subsection formally states the adaptations of the axioms for reference dependent preferences $\{\succsim_r\}_{r \in X}$ in terms of the choice correspondence c . Interpretation is identical to that of those axioms.

Axiom (Category Cancellation*). *For all $x_1, y_1, z_1, x_2, y_2, z_2 \in \mathbb{R}_{++}$ and category k : if $(x_1, z_2) \in c(S^1)$, $(z_1, y_2) \in S^1$, $(z_1, x_2) \in c(S^2)$, $(y_1, z_2) \in S^2$, $(x_1, x_2), (y_1, y_2) \in S^3$ and $S^i \subset K^k(A(S^i))$ for $i \in \{1, 2, 3\}$, then $(x_1, x_2) \in c(S^3)$ whenever $(y_1, y_2) \in c(S^3)$.*

Axiom (Category Monotonicity*). *For any $x, y \in X$: if $x \geq y$ and $x \neq y$, then $(y, k) \not\prec^R (x, k)$ for any category k .*

Axiom (Category Continuity*). *For any $S \in \mathcal{X}$ and any $\epsilon > 0$ so that $E \cap S \setminus c(S) = \emptyset$ where $E \equiv \bigcup_{x \in c(S)} B_\epsilon(x)$ there exists $\delta > 0$ so that if $S' \in \mathcal{X}$, $d(A(S'), A(S)) < \delta$, and for any $y' \in S'$, there is $y \in S$ so that $y' \in B_\delta(y)$, then $c(S') \subset E$ whenever $S' \cap E \neq \emptyset$.*

Define $\succsim^{R,k}$ by $x \succsim^{R,k} y$ if and only if $(x, k) \succsim^R (y, k)$. Using this relation, we can define \oplus^k for each category as we did with preference relations.

Axiom (Affine Across Categories*). *For any $S^1, S^2, S^3 \in \mathcal{X}$, $x^i \in K^j(A(S^i))$, $y^i \in K^k(A(S^i))$ for $i = 1, 2, 3$, and any $\alpha \in (0, 1)$ so that $(x^3, j) \succsim^R (\alpha x^1 \oplus^j (1 - \alpha)x^2, j)$ and $(\alpha y^1 \oplus^k (1 - \alpha)y^2, k) \succsim^R (y^3, k)$: if $x^1 \in c(S^1)$ and $x^2 \in c(S^2)$, then $y^3 \notin c(S^3)$.*

Axiom (Salient Dimension Overvalued*). *For $x, y \in S \cap S'$ with $x_k > y_k$ and $y_{-k} > x_{-k}$, if $x, y \in K^k(A(S))$, $x, y \in R_{-k}(A(S'))$, and $y \in c(S)$, then $x \notin c(S')$.*

Axiom (Reference Interlocking*). *For any $a, b, a', b', x', y', x, y \in X$ with $x_{-i} = a_{-i}$, $y_{-i} = b_{-i}$, $x'_{-i} = a'_{-i}$, $y'_{-i} = b'_{-i}$, $x_i = x'_i$, $y_i = y'_i$, $a_i = a'_i$, $b_i = b'_i$: if $x \sim_k^{R*} y$, $a \succsim_k^{R*} b$, and $x' \sim_j^{R*} y'$, then it does not hold that $b' \succ_j^{R*} a'$.*

B.2. Proof of Theorem 6.

Lemma 12. *Assume that Revealed Structure holds, and that A is a generalized average. If Category-SARP, Category Monotonicity*, Category Cancellation*, and Category Continuity* hold, then for any category k there exists a Category utility U^k so that for any $x, y \in E^{R,k}$,*

$$(x, k) \succsim^R (y, k) \iff U^k(x) \geq U^k(y).$$

Proof. Fix a category i and pick any $x, y \in E^{R,i}$. Let $E^* = E^{R,i} \cap B_{d(x,y)+1}(x)$. As in proof of Lemma 3, there is a continuous path $\theta : [0, 1] \rightarrow E^*$ so that $\theta(0) = x$ and $\theta(1) = y$ that crosses each $\succsim^{R,i}$ indifference curve at most once, and $Y = \theta^{-1}([0, 1])$ is compact. We will show that for any $z \in Y$, there exists an open set $z \in B_z \subset E^*$ so that $\succsim^{R,i}$ is complete on B_z . If this is the case, we can mimic the rest of the proof of Lemma 3 to show that either $x \succsim^{R,i} y$ or $y \succsim^{R,i} x$.

By definition of E^* , for any $z \in E^*$, there exists $S \in \mathcal{X}$ with $A(S) = r$ so that $c(S) = z$. Since $K^i(r)$ is open, there exists $\epsilon_1 > 0$ so that $B_{2\epsilon_1}(z) \subset K^i(r)$. By Lemma 11, there exists $\epsilon_2 > 0$ so that $r' \in B_{\epsilon_2}(r)$ implies $B_{\epsilon_1}(z) \subset K^i(r')$. Pick $\zeta \in (0, \frac{1}{2})$ so that $B_\zeta(z) \cap S = z$. By Category Continuity*, there exists $\epsilon_3 > 0$ so that for any $S' \in \mathcal{X}$ with $d(A(S'), A(S)) < \epsilon_3$, for any $y' \in S'$, there is $y \in S$ so that $y' \in B_{\epsilon_3}(y)$, and $S' \cap B_\zeta(x) \neq \emptyset$, then $c(S') \subset B_\zeta(x)$. By Generalized Average, there exists $\epsilon_4 > 0$ so that $z' \in B_{\epsilon_4}(z)$ implies $d(A(S \setminus \{z\} \cup \{z'\}), A(S)) < \min\{\epsilon_2, \epsilon_3\}/2$. Let $\epsilon^* = \min\{\epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4, \zeta\}$.

Pick any $x', y' \in B_{\epsilon^*/4}(z)$ and let $z_* = z - \frac{1}{2}\epsilon^*$. Set $S_0 = S \setminus \{z\} \cup \{z_*\}$, noting $d(r, A(S_0)) < \epsilon_2/2$. By Generalized Average, there exists S^* with $\{x', y'\} \cup S_0 \subset S^*$ so that $d(A(S^*), A(S_0)) < \epsilon^*/2$ and $d(S_0, S^* \setminus [\{x', y'\} \cup S_0]) < (\epsilon^*/2)^2$. Since $d(A(S^*), r) \leq d(A(S^*), A(S_0)) + d(A(S_0), r) < \epsilon_2$, $x', y' \in K^i(A(S^*))$. Since every member of S^* is no more than ϵ^* away from a member of S , Category Continuity* implies that $c(S^*) \subset B_\zeta(z)$. CM* gives that either $x' \in c(S^*)$ or $y' \in c(S^*)$, so $x' \succsim^{R,i} y'$ or $y' \succsim^{R,i} x'$.

Continuity follows along the same lines as Lemma 2. CM* gives that it is also monotone, and Category Cancellation* that it is locally additive. Apply Theorem 2.2 of Chateauneuf & Wakker [1993] to get a globally additive representation U^k . \square

By Lemma 12, there exists a category utility U^k for each category. Since $E^{R,k}$ is dense in D^k , we can extend U^k to D^k uniquely. By Generalized Average and Category Continuity*, for any $S \in X$ with $z \in [D^k \setminus E^{R,k}] \cap S$, there is a $z' \in E^{R,k}$ arbitrarily close to z so that $c(S) = c([S \setminus \{z\}] \cup \{z'\})$, so it is sufficient to establish a representation when all alternatives categorized as k in S belong $E^{R,k}$ for each k and S .

Fix two regions k and j . By CAR, for any $x \in E^{R,k}$ there exists $x' \in E^{R,k}$, $y \in E^{R,j}$, and $S \in \mathcal{X}$ so that $x', y \in c(S)$ and $x \sim^{R,k} x'$. This implies there exists a strictly increasing function H so that $V(x|r) = U^k(x)$ when $x \in K^k(r)$ and $V(x|r) = H(U^j(x))$ when $x \in K^j(r)$ represents choice (when $S \subset K^k \cup K^j$). This is well-defined and represents choice by Category SARP. By AAC*, H is an affine function. The argument are readily seen to extend inductively to all regions, which complete the proof. \square

B.3. Proof of Lemma 1. Pick any $x \in X$ and set $S = \{x, x'\}$ where $x' = (\frac{1}{2}x_1, x_2)$. Then, $A(S)_2 = x_2$ by strong generalized average, so both x and x' are 1-salient by S4. By CM*, $x \in c(S)$, and so $x \in E^{R,1}$. x was arbitrary, so $X = E^{R,1}$. Similar for K^2 . \square

B.4. Proof of Proposition 3. By Lemma 1, the structure assumption is satisfied. By Theorem 5, the category function is generated by a salience function. By Theorem 6, c conforms to Strong CTM. Mimicking the arguments of Theorem 2, Reference Interlocking implies $U^k(x) = w_1^k u_1(x_1) + w_2^k u_2(x_2) + \beta_k$. The rest follows from the arguments that establish Proposition 2. \square

B.5. Proof of Proposition 4. Pick any r , and suppose $U^k(r) \geq U^{-k}(r)$. Since A is a generalized average, for any $y \gg r$ there exists a menu S so that $A(S)$ is arbitrarily close to r and $y \gg x'$ for all $x' \in S$ (pick S so its convex hull is in a small enough neighborhood of r that doesn't include y). By making that neighborhood smaller if necessary, either y belongs to $K^k(A(S))$ or $y \notin K^k(r)$. There exists a y' arbitrarily close to, but not equal to, y , so that $U^k(y') = U^k(y)$ and $U^{-k}(y) \neq U^{-k}(y')$. In the former case either y or y' is chosen from the menu S' by categorical monotonicity*, where S' is a menu (assumed to exist by generalized average) with $A(S')$ sufficiently close to $A(S)$ and $y, y' \in S'$. Moreover, $y' \in c(S')$ if it is close enough that it too belongs to K^k . Conclude that $y \in K^k(r)$ implies that there exists y' arbitrarily close to, but not equal to, y with $U^k(y') = U^k(y)$ and $\{y, y'\} = c(S')$. If $y \notin K^k(r)$, then both y and y' cannot be chosen. Either y is not chosen because it is in K^{-k} , or the DM will not be indifferent between y' and y . The rest follows from Proposition 1. \square